




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/82, 15/54, C08B 30/04, A01H 5/00</b>		A2	(11) International Publication Number: <b>WO 96/34968</b>
(21) International Application Number: <b>PCT/GB96/01075</b>		(43) International Publication Date: <b>7 November 1996 (07.11.96)</b>	
(22) International Filing Date: <b>3 May 1996 (03.05.96)</b>		(74) Agent: <b>KEITH W. NASH &amp; CO.; 90-92 Regent Street, Cambridge CB2 1DP (GB).</b>	
(30) Priority Data: 9509229.2      5 May 1995 (05.05.95)      GB 9607409.1      10 April 1996 (10.04.96)      GB		(81) Designated States: <b>AU, BR, CA, JP, KR, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</b>	
(71) Applicant (for all designated States except US): <b>NATIONAL STARCH AND CHEMICAL INVESTMENT HOLDING CORPORATION [US/US]; Suite 27, 501 Silverside Road, Wilmington, DE 19809 (US).</b>		Published Without international search report and to be republished upon receipt of that report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>COOKE, David [GB/GB]; 2 Hunts Path, Oakley, Bedfordshire MK43 7SR (GB). DEBET, Martine [FR/GB]; 12 Bentley Close, Rectory Farm, Northampton NN3 5JS (GB). GIDLEY, Michael, John [GB/GB]; 45 Holmes Avenue, Raunds, Northamptonshire NN9 6SZ (GB). JOBLING, Stephen, Alan [GB/GB]; 19 Burwell Road, Eaton Socon, Huntingdon PE19 3QQ (GB). SAFFORD, Richard [GB/GB]; 10 Furness Close, Bedford, Bedfordshire MK41 8RN (GB). SIDEBOTTOM, Christopher, Michael [GB/GB]; 35 Waterloo Road, Bedford, Bedfordshire MK40 3PQ (GB). WESTCOTT, Roger, John</b>			

(54) Title: IMPROVEMENTS IN OR RELATING TO PLANT STARCH COMPOSITION

BEST AVAILABLE COPY



## (57) Abstract

Disclosed is a nucleotide sequence encoding an effective portion of a class A starch branching enzyme (SBE) obtainable from potato plants, or a functional equivalent thereof, together with, inter alia, a corresponding polypeptide, a method of altering the characteristics of a plant, a plant having altered characteristics; and starch, particularly starch obtained from a potato plant, having novel properties.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**Title:** Improvements in or Relating to Plant Starch Composition

### **Field of the Invention**

This invention relates to novel nucleotide sequences, polypeptides encoded thereby, vectors and host cells and host organisms comprising one or more of the novel sequences, and to a method of altering one or more characteristics of an organism. The invention also relates to starch having novel properties and to uses thereof.

### **Background of the Invention**

Starch is the major form of carbon reserve in plants, constituting 50% or more of the dry weight of many storage organs - e.g. tubers, seeds of cereals. Starch is used in numerous food and industrial applications. In many cases, however, it is necessary to modify the native starches, via chemical or physical means, in order to produce distinct properties to suit particular applications. It would be highly desirable to be able to produce starches with the required properties directly in the plant, thereby removing the need for additional modification. To achieve this via genetic engineering requires knowledge of the metabolic pathway of starch biosynthesis. This includes characterisation of genes and encoded gene products which catalyse the synthesis of starch. Knowledge about the regulation of starch biosynthesis raises the possibility of "re-programming" biosynthetic pathways to create starches with novel properties that could have new commercial applications.

The commercially useful properties of starch derive from the ability of the native granular form to swell and absorb water upon suitable treatment. Usually heat is required to cause granules to swell in a process known as gelatinisation, which has been defined (W A Atwell *et al*, Cereal Foods World 33, 306-311, 1988) as "*... the collapse (disruption) of molecular orders within the starch granule manifested in irreversible changes in properties such as granular swelling, native crystallite melting, loss of birefringence, and starch solubilisation. The point of initial gelatinisation and the range over which it occurs is governed by starch concentration, method of observation, granule type, and heterogeneities within the granule population under observation*". A number of techniques are available

for the determination of gelatinisation as induced by heating, a convenient and accurate method being differential scanning calorimetry, which detects the temperature range and enthalpy associated with the collapse of molecular orders within the granule. To obtain accurate and meaningful results, the peak and/or onset temperature of the endotherm observed by differential scanning calorimetry is usually determined.

The consequence of the collapse of molecular orders within starch granules is that the granules are capable of taking up water in a process known as pasting, which has been defined (W A Atwell *et al*, Cereal Foods World 33, 306-311, 1988) as "*... the phenomenon following gelatinisation in the dissolution of starch. It involves granular swelling, exudation of molecular components from the granule, and eventually, total disruption of the granules*". The best method of evaluating pasting properties is considered to be the viscoamylograph (Atwell *et al*, 1988 cited above) in which the viscosity of a stirred starch suspension is monitored under a defined time/temperature regime. A typical viscoamylograph profile for potato starch shows an initial rise in viscosity, which is considered to be due to granule swelling. In addition to the overall shape of the viscosity response in a viscoamylograph, a convenient quantitative measure is the temperature of initial viscosity development (onset). Figure 1 shows such a typical viscosity profile for potato starch, during and after cooking, and includes stages A-D which correspond to viscosity onset (A), maximum viscosity (B), complete dispersion (C) and reassociation of molecules (or retrogradation, D). In the figure, the dotted line represents viscosity (in stirring number units) of a 10% w/w starch suspension and the unbroken line shows the temperature in degrees centigrade. At a certain point, defined by the viscosity peak, granule swelling is so extensive that the resulting highly expanded structures are susceptible to mechanically-induced fragmentation under the stirring conditions used. With increased heating and holding at 95°C, further reduction in viscosity is observed due to increased fragmentation of swollen granules. This general profile has previously always been found for native potato starch.

After heating starches in water to 95°C and holding at that temperature (for typically 15 minutes), subsequent cooling to 50°C results in an increase in viscosity due to the process of retrogradation or set-back. Retrogradation (or set-back) is defined (Atwell *et al*., 1988

cited above) as "...a process which occurs when the molecules comprising gelatinised starch begin to reassociate in an ordered structure...". At 50°C, it is primarily the amylose component which reassociates, as indicated by the increase in viscoamylograph viscosity for starch from normal maize (21.6% amylose) compared with starch from waxy maize (1.1% amylose) as shown in Figure 2. Figure 2 is a viscoamylograph of 10%w/w starch suspensions from waxy maize (solid line), conventional maize (dots and dashes), high amylose variety (hylon 5, dotted line) and a very high amylose variety (hylon 7, crosses). The temperature profile is also shown by a solid line, as in Figure 1. The extent of viscosity increase in the viscoamylograph on cooling and holding at 50°C depends on the amount of amylose which is able to reassociate due to its exudation from starch granules during the gelatinisation and pasting processes. A characteristic of amylose-rich starches from maize plants is that very little amylose is exuded from granules by gelatinisation and pasting up to 95°C, probably due to the restricted swelling of the granules. This is illustrated in Figure 2 which shows low viscosities for a high amylose (44.9%) starch (Hylon 5) from maize during gelatinisation and pasting at 95°C and little increase in viscosity on cooling and holding at 50°C. This effect is more extreme for a higher amylose content (58%, as in Hylon 7), which shows even lower viscosities in the viscoamylograph test (Figure 2). For commercially-available high amylose starches (currently available from maize plants, such as those described above), processing at greater than 100°C is usually necessary in order to generate the benefits of high amylose contents with respect to increased rates and strengths of reassociation, but use of such high temperatures is energetically unfavourable and costly. Accordingly, there is an unmet need for starches of high amylose content which can be processed below 100°C and still show enhanced levels of reassociation, as indicated for example by viscoamylograph measurements.

The properties of potato starch are useful in a variety of both food and non-food (paper, textiles, adhesives etc.) applications. However, for many applications, properties are not optimum and various chemical and physical modifications well known in the art are undertaken in order to improve useful properties. Two types of property manipulation which would be of use are: the controlled alteration of gelatinisation and pasting temperatures; and starches which suffer less granular fragmentation during pasting than

conventional starches.

Currently the only ways of manipulating the gelatinisation and pasting temperatures of potato starch are by the inclusion of additives such as sugars, polyhydroxy compounds of salts (Evans & Haisman, *Starke* 34, 224-231, 1982) or by extensive physical or chemical pre-treatments (e.g. Stute, *Starke* 44, 205-214, 1992). The reduction of granule fragmentation during pasting can be achieved either by extensive physical pretreatments (Stute, *Starke* 44, 205-214, 1992) or by chemical cross-linking. Such processes are inconvenient and inefficient. It is therefore desirable to obtain plants which produce starch which intrinsically possesses such advantageous properties.

Starch consists of two main polysaccharides, amylose and amylopectin. Amylose is a generally linear polymer containing  $\alpha$ -1,4 linked glucose units, while amylopectin is a highly branched polymer consisting of a  $\alpha$ -1,4 linked glucan backbone with  $\alpha$ -1,6 linked glucan branches. In most plant storage reserves amylopectin constitutes about 75% of the starch content. Amylopectin is synthesized by the concerted action of soluble starch synthase and starch branching enzyme [ $\alpha$ -1,4 glucan:  $\alpha$ -1,4 glucan 6-glycosyltransferase, EC 2.4.1.18]. Starch branching enzyme (SBE) hydrolyses  $\alpha$ -1,4 linkages and rejoins the cleaved glucan, via an  $\alpha$ -1,6 linkage, to an acceptor chain to produce a branched structure. The physical properties of starch are strongly affected by the relative abundance of amylose and amylopectin, and SBE is therefore a crucial enzyme in determining both the quantity and quality of starches produced in plant systems.

In most plants studied to date e.g. maize (Boyer & Preiss, 1978 *Biochem. Biophys. Res. Comm.* 80, 169-175), rice (Smyth, 1988 *Plant Sci.* 57, 1-8) and pea (Smith, *Planta* 175, 270-279), two forms of SBE have been identified, each encoded by a separate gene. A recent review by Burton *et al.*, (1995 *The Plant Journal* 7, 3-15) has demonstrated that the two forms of SBE constitute distinct classes of the enzyme such that, in general, enzymes of the same class from different plants may exhibit greater similarity than enzymes of different classes from the same plant. In their review, Burton *et al.* termed the two respective enzyme families class "A" and class "B", and the reader is referred thereto (and to the references cited therein) for a detailed discussion of the distinctions

between the two classes. One general distinction of note would appear to be the presence, in class A SBE molecules, of a flexible N-terminal domain, which is not found in class B molecules. The distinctions noted by Burton *et al.* are relied on herein to define class A and class B SBE molecules, which terms are to be interpreted accordingly.

However in potato, only one isoform of the SBE molecule (belonging to class B) has thus far been reported and only one gene cloned (Blennow & Johansson, 1991 *Phytochem.* 30, 437-444, and Koßmann *et al.*, 1991 *Mol. Gen. Genet.* 230, 39-44). Further, published attempts to modify the properties of starch in potato plants (by preventing expression of the single known SBE) have generally not succeeded (e.g. Müller-Rober & Koßmann 1994 *Plant Cell and Environment* 17, 601-613).

### Summary of the Invention

In a first aspect the invention provides a nucleotide sequence encoding an effective portion of a class A starch branching enzyme (SBE) obtainable from potato plants.

Preferably the nucleotide sequence encodes a polypeptide comprising an effective portion of the amino acid sequence shown in Figure 5 (excluding the sequence MNKRIDL, which does not represent part of the SBE molecule), or a functional equivalent thereof (which term is discussed below). The amino acid sequence shown in Figure 5 (Seq ID No. 15) includes a leader sequence which directs the polypeptide, when synthesised in potato cells, to the amyloplast. Those skilled in the art will recognise that the leader sequence is removed to produce a mature enzyme and that the leader sequence is therefore not essential for enzyme activity. Accordingly, an "effective portion" of the polypeptide is one which possesses sufficient SBE activity to complement the branching enzyme mutation in *E. coli* KV 832 cells (described below) and which is active when expressed in *E. coli* in the phosphorylation stimulation assay. An example of an incomplete polypeptide which nevertheless constitutes an "effective portion" is the mature enzyme lacking the leader sequence. By analogy with the pea class A SBE sequence, the potato class A sequence shown in Figure 5 probably possesses a leader sequence of about 48 amino acid residues, such that the N terminal amino acid sequence is thought to commence around the glutamic acid residue (E) at position 49 (EKSSYN... etc.). Those skilled in the art will appreciate

that an effective portion of the enzyme may well omit other parts of the sequence shown in the figure without substantial detrimental effect. For example, the C-terminal glutamic acid-rich region could be reduced in length, or possibly deleted entirely, without abolishing class A SBE activity. A comparison with other known SBE sequences, especially other class A SBE sequences (see for example, Burton *et al.*, 1995 cited above), should indicate those portions which are highly conserved (and thus likely to be essential for activity) and those portions which are less well conserved (and thus are more likely to tolerate sequence changes without substantial loss of enzyme activity).

Conveniently the nucleotide sequence will comprise substantially nucleotides 289 to 2790 of the DNA sequence (Seq ID No. 14) shown in Figure 5 (which nucleotides encode the mature enzyme) or a functional equivalent thereof, and may also include further nucleotides at the 5' or 3' end. For example, for ease of expression, the sequence will desirably also comprise an in-frame ATG start codon, and may also encode a leader sequence. Thus, in one embodiment, the sequence further comprises nucleotides 145 to 288 of the sequence shown in Figure 5. Other embodiments are nucleotides 228 to 2855 of the sequence labelled "psbe2con.seq" in Figure 8, and nucleotides 57 to 2564 of the sequence shown in Figure 12 (preferably comprising an in-frame ATG start codon, such as the sequence of nucleotides 24 to 56 in the same Figure), or functional equivalents of the aforesaid sequences.

The term "functional equivalent" as applied herein to nucleotide sequences is intended to encompass those sequences which differ in their nucleotide composition to that shown in Figure 5 but which, by virtue of the degeneracy of the genetic code, encode polypeptides having identical or substantially identical amino acid sequences. It is intended that the term should also apply to sequences which are sufficiently homologous to the sequence of the invention that they can hybridise to the complement thereof under stringent hybridisation conditions - such equivalents will preferably possess at least 85%, more preferably at least 90%, and most preferably at least 95% sequence homology with the sequence of the invention as exemplified by nucleotides 289 to 2790 of the DNA sequence shown in Figure 5. It will be apparent to those skilled in the art that the nucleotide sequence of the invention may also find useful application when present as an "antisense"



sequence. Accordingly, functionally equivalent sequences will also include those sequences which can hybridise, under stringent hybridisation conditions, to the sequence of the invention (rather than the complement thereof). Such "antisense" equivalents will preferably possess at least 85%, more preferably at least 90%, and most preferably 95% sequence homology with the complement of the sequence of the invention as exemplified by nucleotides 289 to 2790 of the DNA sequence shown in Figure 5. Particular functional equivalents are shown, for example, in Figures 8 and 10 (if one disregards the various frameshift mutations noted therein).

The invention also provides vectors, particularly expression vectors, comprising the nucleotide sequence of the invention. The vector will typically comprise a promoter and one or more regulatory signals of the type well known to those skilled in the art. The invention also includes provision of cells transformed (which term encompasses transduction and transfection) with a vector comprising the nucleotide sequence of the invention.

The invention further provides a class A SBE polypeptide, obtainable from potato plants. In particular the invention provides the polypeptide in substantially pure form, especially in a form free from other plant-derived (especially potato plant-derived) components, which can be readily accomplished by expression of the relevant nucleotide sequence in a suitable non-plant host (such as any one of the yeast strains routinely used for expression purposes, e.g. *Pichia spp.* or *Saccharomyces spp.*). Typically the enzyme will substantially comprise the sequence of amino acid residues 49 to 882 shown in Figure 5 (disregarding the sequence MNKRIDL, which is not part of the enzyme), or a functional equivalent thereof. The polypeptide of the invention may be used in a method of modifying starch *in vitro*, comprising treating starch under suitable conditions (e.g. appropriate temperature, pH, etc) with an effective amount of the polypeptide according to the invention.

The term "functional equivalent", as applied herein to amino acid sequences, is intended to encompass amino acid sequences substantially similar to that shown in Figure 5, such that the polypeptide possesses sufficient activity to complement the branching enzyme mutation in *E. coli* KV 832 cells (described below) and which is active in *E. coli* in the

phosphorylation stimulation assay. Typically such functionally equivalent amino acid sequences will preferably possess at least 85%, more preferably at least 90%, and most preferably at least 95% sequence identity with the amino acid sequence of the mature enzyme (i.e. minus leader sequence) shown in Figure 5. Those skilled in the art will appreciate that conservative substitutions may be made generally throughout the molecule without substantially affecting the activity of the enzyme. Moreover, some non-conservative substitutions may be tolerated, especially in the less highly conserved regions of the molecule. Such substitutions may be made, for example, to modify slightly the activity of the enzyme. The polypeptide may, if desired, include a leader sequence, such as that exemplified by residues 1 to 48 of the amino acid sequence shown in Figure 5, although other leader sequences and signal peptides and the like are known and may be included.

A portion of the nucleotide sequence of the invention has been introduced into a plant and found to affect the characteristics of the plant. In particular, introduction of the sequence of the invention, operably linked in the antisense orientation to a suitable promoter, was found to reduce the amount of branched starch molecules in the plant. Additionally, it has recently been demonstrated in other experimental systems that "sense suppression" can also occur (i.e. expression of an introduced sequence operably linked in the sense orientation can interfere, by some unknown mechanism, with the expression of the native gene), as described by Matzke & Matzke (1995 *Plant Physiol.* 107, 679-685). Any one of the methods mentioned by Matzke & Matzke could, in theory, be used to affect the expression in a host of a homologous SBE gene.

It is believed that antisense methods are mainly operable by the production of antisense mRNA which hybridises to the sense mRNA, preventing its translation into functional polypeptide, possibly by causing the hybrid RNA to be degraded (e.g. Sheehy *et al.*, 1988 *PNAS* 85, 8805-8809; Van der Krol *et al.*, *Mol. Gen. Genet.* 220, 204-212). Sense suppression also requires homology between the introduced sequence and the target gene, but the exact mechanism is unclear. It is apparent however that, in relation to both antisense and sense suppression, neither a full length nucleotide sequence, nor a "native" sequence is essential. Preferably the "effective portion" used in the method will comprise

at least one third of the full length sequence, but by simple trial and error other fragments (smaller or larger) may be found which are functional in altering the characteristics of the plant.

Thus, in a further aspect the invention provides a method of altering the characteristics of a plant, comprising introducing into the plant an effective portion of the sequence of the invention operably linked to a suitable promoter active in the plant. Conveniently the sequence will be linked in the anti-sense orientation to the promoter. Preferably the plant is a potato plant. Conveniently, the characteristic altered relates to the starch content and/or starch composition of the plant (i.e. amount and/or type of starch present in the plant). Preferably the method of altering the characteristics of the plant will also comprise the introduction of one or more further sequences, in addition to an effective portion of the sequence of the invention. The introduced sequence of the invention and the one or more further sequences (which may be sense or antisense sequences) may be operably linked to a single promoter (which would ensure both sequences were transcribed at essentially the same time), or may be operably linked to separate promoters (which may be necessary for optimal expression). Where separate promoters are employed they may be identical to each other or different. Suitable promoters are well known to those skilled in the art and include both constitutive and inducible types. Examples include the CaMV 35S promoter (e.g. single or tandem repeat) and the patatin promoter. Advantageously the promoter will be tissue-specific. Desirably the promoter will cause expression of the operably linked sequence at substantial levels only in the tissue of the plant where starch synthesis and/or starch storage mainly occurs. Thus, for example, where the sequence is introduced into a potato plant, the operably linked promoter may be tuber-specific, such as the patatin promoter.

Desirably, for example, the method will also comprise the introduction of an effective portion of a sequence encoding a class B SBE, operably linked in the antisense orientation to a suitable promoter active in the plant. Desirably the further sequence will comprise an effective portion of the sequence encoding the potato class B SBE molecule. Conveniently the further sequence will comprise an effective portion of the sequence described by Blennow & Johansson (1991 *Phytochem.* 30, 437-444) or that disclosed in

WO92/11375. More preferably, the further sequence will comprise at least an effective portion of the sequence disclosed in International Patent Application No. WO 95/26407. Use of antisense sequences against both class A and class B SBE in combination has now been found by the present inventors to result in the production of starch having very greatly altered properties (see below). Those skilled in the art will appreciate the possibility that, if the plant already comprises a sense or antisense sequence which efficiently inhibits the class B SBE activity, introduction of a sense or antisense sequence to inhibit class A SBE activity (thereby producing a plant with inhibition of both class A and class B activity) might alter greatly the properties of the starch in the plant, without the need for introduction of one or more further sequences. Thus the sequence of the invention is conveniently introduced into plants already having low levels of class A and/or class B SBE activity, such that the inhibition resulting from the introduction of the sequence of the invention is likely to have a more pronounced effect.

The sequence of the invention, and the one or more further sequences if desired, can be introduced into the plant by any one of a number of well-known techniques (e.g. Agrobacterium-mediated transformation, or by "biolistic" methods). The sequences are likely to be most effective in inhibiting SBE activity in potato plants, but theoretically could be introduced into any plant. Desirable examples include pea, tomato, maize, wheat, rice, barley, sweet potato and cassava plants. Preferably the plant will comprise a natural gene encoding an SBE molecule which exhibits reasonable homology with the introduced nucleic acid sequence of the invention.

In another aspect, the invention provides a plant cell, or a plant or the progeny thereof, which has been altered by the method defined above. The progeny of the altered plant may be obtained, for example, by vegetative propagation, or by crossing the altered plant and reserving the seed so obtained. The invention also provides parts of the altered plant, such as storage organs. Conveniently, for example, the invention provides tubers comprising altered starch, said tubers being obtained from an altered plant or the progeny thereof. Potato tubers obtained from altered plants (or the progeny thereof) will be particularly useful materials in certain industrial applications and for the preparation and/or processing of foodstuffs and may be used, for example, to prepare low-fat waffles and

chips (amylose generally being used as a coating to prevent fat uptake), and to prepare mashed potato (especially "instant" mashed potato) having particular characteristics.

In particular relation to potato plants, the invention provides a potato plant or part thereof which, in its wild type possesses an effective SBE A gene, but which plant has been altered such that there is no effective expression of an SBE A polypeptide within the cells of at least part of the plant. The plant may have been altered by the method defined above, or may have been selected by conventional breeding to be deleted for the class A SBE gene, presence or absence of which can be readily determined by screening samples of the plants with a nucleic acid probe or antibody specific for the potato class A gene or gene product respectively.

The invention also provides starch extracted from a plant altered by the method defined above, or the progeny of such a plant, the starch having altered properties compared to starch extracted from equivalent, but unaltered, plants. The invention further provides a method of making altered starch, comprising altering a plant by the method defined above and extracting therefrom starch having altered properties compared to starch extracted from equivalent, but unaltered, plants. Use of nucleotide sequences in accordance with the invention has allowed the present inventors to produce potato starches having a wide variety of novel properties.

In particular the invention provides the following: a plant (especially a potato plant) altered by the method defined above, containing starch which, when extracted from the plant, has an elevated endotherm peak temperature as judged by DSC, compared to starch extracted from a similar, but unaltered, plant; a plant (especially a potato plant) altered by the method defined above, containing starch which, when extracted from the plant, has an elevated viscosity onset temperature (conveniently elevated by 10 - 25°C) as judged by viscoamylograph compared to starch extracted from a similar, but unaltered, plant; a plant (especially a potato plant) altered by the method defined above, containing starch which, when extracted from the plant, has a decreased peak viscosity (conveniently decreased by 240 - 700SNUs) as judged by viscoamylograph compared to starch extracted from a similar, but unaltered, plant; a plant (especially a potato plant) altered by the method

defined above, containing starch which, when extracted from the plant, has an increased pasting viscosity (conveniently increased by 37 - 260 SNUs) as judged by viscoamylograph compared to starch extracted from a similar, but unaltered, plant; a plant (especially a potato plant) altered by the method defined above, containing starch which, when extracted from the plant, has an increased set-back viscosity (conveniently increased by 224 - 313 SNUs) as judged by viscoamylograph compared to starch extracted from a similar, but unaltered, plant; a plant (especially a potato plant) altered by the method defined above, containing starch which, when extracted from the plant, has a decreased set-back viscosity as judged by viscoamylograph compared to starch extracted from a similar, but unaltered, plant; and a plant (especially a potato plant) altered by the method defined above, containing starch which, when extracted from the plant, has an elevated amylose content as judged by iodometric assay (i.e. by the method of Morrison & Laignelet 1983, cited above) compared to starch extracted from a similar, but unaltered, plant. The invention also provides for starch obtainable or obtained from such plants as aforesaid.

In particular the invention provides for starch which, as extracted from a potato plant by wet milling at ambient temperature, has one or more of the following properties, as judged by viscoamylograph analysis performed according to the conditions defined below: viscosity onset temperature in the range 70-95°C (preferably 75-95°C); peak viscosity in the range 500 - 12 stirring number units; pasting viscosity in the range 214 - 434 stirring number units; set-back viscosity in the range 450 - 618 or 14 - 192 stirring number units; or displays no significant increase in viscosity during viscoamylograph. Peak, pasting and set-back viscosities are defined below. Viscosity onset temperature is the temperature at which there is a sudden, marked increase in viscosity from baseline levels during viscoamylograph, and is a term well-known to those skilled in the art.

In other particular embodiments, the invention provides starch which as extracted from a potato plant by wet milling at ambient temperature has a peak viscosity in the range 200 - 500 SNUs and a set-back viscosity in the range 275-618 SNUs as judged by viscoamylograph according to the protocol defined below; and starch which as extracted from a potato plant by wet milling at ambient temperature has a viscosity which does not decrease between the start of the heating phase (step 2) and the start of the final holding

phase (step 5) and has a set-back viscosity of 303 SNU's or less as judged by viscoamylograph according to the protocol defined below.

For the purposes of the present invention, viscoamylograph conditions are understood to pertain to analysis of a 10% (w/w) aqueous suspension of starch at atmospheric pressure, using a Newport Scientific Rapid Visco Analyser with a heating profile of: holding at 50°C for 2 minutes (step 1), heating from 50 to 95°C at a rate of 1.5°C per minute (step 2), holding at 95°C for 15 minutes (step 3), cooling from 95 to 50°C at a rate of 1.5°C per minute (step 4), and then holding at 50°C for 15 minutes (step 5). Peak viscosity may be defined for present purposes as the maximum viscosity attained during the heating phase (step 2) or the holding phase (step 3) of the viscoamylograph. Pasting viscosity may be defined as the viscosity attained by the starch suspensions at the end of the holding phase (step 3) of the viscoamylograph. Set-back viscosity may be defined as the viscosity of the starch suspension at the end of step 5 of the viscoamylograph.

In yet another aspect the invention provides starch from a potato plant having an apparent amylose content (% w/w) of at least 35%, as judged by iodometric assay according to the method described by Morrison & Laignelet (1983 *J. Cereal Science* 1, 9-20). Preferably the starch will have an amylose content of at least 40%, more preferably at least 50%, and most preferably at least 66%. Starch obtained directly from a potato plant and having such properties has not hitherto been produced. Indeed, as a result of the present invention, it is now possible to generate *in vivo* potato starch which has some properties analogous to the very high amylose starches (e.g. Hylon 7) obtainable from maize.

Starches with high (at least 35%) amylose contents find commercial application as, amongst other reasons, the amylose component of starch reassociates more strongly and rapidly than the amylopectin component during retrogradation processes. This may result, for example, in pastes with higher viscosities, gels of greater cohesion, or films of greater strength for starches with high (at least 35%) compared with normal (less than 35%) amylose contents. Alternatively, starches may be obtained with very high amylose contents, such that the granule structure is substantially preserved during heating, resulting in starch suspensions which demonstrate substantially no increase in viscosity during

cooking (i.e. there is no significant viscosity increase during viscoamylograph conditions defined above). Such starches typically exhibit a viscosity increase of less than 10% (preferably less than 5%) during viscoamylograph under the conditions defined above.

In commerce, these valuable properties are currently obtained from starches of high amylose content derived from maize plants. It would be of commercial value to have an alternative source of high amylose starches from potato as other characteristics such as granule size, organoleptic properties and textural qualities may distinguish application performances of high amylose starches from maize and potato plants.

Thus high amylose starch obtained by the method of the present invention may find application in many different technological fields, which may be broadly categorised into two groups: food products and processing; and "Industrial" applications. Under the heading of food products, the novel starches of the present invention may find application as, for example, films, barriers, coatings or gelling agents. In general, high amylose content starches absorb less fat during frying than starches with low amylose content, thus the high amylose content starches of the invention may be advantageously used in preparing low fat fried products (e.g. potato chips, crisps and the like). The novel starches may also be employed with advantage in preparing confectionery and in granular and retrograded "resistant" starches. "Resistant" starch is starch which is resistant to digestion by  $\alpha$ -amylase. As such, resistant starch is not digested by  $\alpha$ -amylases present in the human small intestine, but passes into the colon where it exhibits properties similar to soluble and insoluble dietary fibre. Resistant starch is thus of great benefit in foodstuffs due to its low calorific value and its high dietary fibre content. Resistant starch is formed by the retrogradation (akin to recrystallization) of amylose from starch gels. Such retrogradation is inhibited by amylopectin. Accordingly, the high amylose starches of the present invention are excellent starting materials for the preparation of resistant starch. Suitable methods for the preparation of resistant starch are well-known to those skilled in the art and include, for example, those described in US 5,051,271 and US 5,281,276. Conveniently the resistant starches provided by the present invention comprise at least 5% total dietary fibre, as judged by the method of Prosky *et al.*, (1985 J. Assoc. Off. Anal. Chem. 68, 677), mentioned in US 5,281, 276.



Under the heading of "Industrial" applications, the novel starches of the invention may be advantageously employed, for example, in corrugating adhesives, in biodegradable products such as loose fill packaging and foamed shapes, and in the production of glass fibers and textiles.

Those skilled in the art will appreciate that the novel starches of the invention may, if desired, be subjected *in vitro* to conventional enzymatic, physical and/or chemical modification, such as cross-linking, introduction of hydrophobic groups (e.g. octenyl succinic acid, dodecyl succinic acid), or derivatization (e.g. by means of esterification or etherification).

In yet another aspect the invention provides high (35% or more) amylose starches which generate paste viscosities greater than those obtained from high amylose starches from maize plants after processing at temperatures below 100°C. This provides the advantage of more economical starch gelatinisation and pasting treatments through the use of lower processing temperatures than are currently required for high amylose starches from maize plants.

The invention will now be further described by way of illustrative example and with reference to the drawings, of which:

Figure 1 shows a typical viscoamylograph for a 10% w/w suspension of potato starch;

Figure 2 shows viscoamylographs for 10% suspensions of starch from various maize varieties;

Figure 3 is a schematic representation of the cloning strategy used by the present inventors;

Figure 4a shows the amino acid alignment of the C-terminal portion of starch branching enzyme isoforms from various sources: amino acid residues matching the consensus

sequence are shaded;

Figure 4b shows the alignment of DNA sequences of various starch branching enzyme isoforms which encode a conserved amino acid sequence;

Figure 5 shows the DNA sequence (Seq ID No. 14) and predicted amino acid sequence (Seq ID No. 15) of a full length potato class A SBE cDNA clone obtained by PCR;

Figure 6 shows a comparison of the most highly conserved part of the amino acid sequences of potato class A (uppermost sequence) and class B (lowermost sequence) SBE molecules;

Figure 7 shows a comparison of the amino acid sequence of the full length potato class A (uppermost sequence) and pea (lowermost sequence) class A SBE molecules;

Figure 8 shows a DNA alignment of various full length potato class A SBE clones obtained by the inventors;

Figure 9 shows the DNA sequence of a potato class A SBE clone determined by direct sequencing of PCR products, together with the predicted amino acid sequence;

Figure 10 is a multiple DNA alignment of various full length potato SBE A clones obtained by the inventors;

Figure 11 is a schematic illustration of the plasmid pSJ64;

Figure 12 shows the DNA sequence and predicted amino acid sequence of the full length potato class A SBE clone as present in the plasmid pSJ90; and

Figure 13 shows viscoamylographs for 10% w/w suspensions of starch from various transgenic potato plants made by the relevant method aspect of the invention.

## Examples

### Example 1

#### Cloning of Potato class A SBE

The strategy for cloning the second form of starch branching enzyme from potato is shown in Figure 3. The small arrowheads represent primers used by the inventors in PCR and RACE protocols. The approximate size of the fragments isolated is indicated by the numerals on the right of the Figure. By way of explanation, a comparison of the amino acid sequences of several cloned plant starch branching enzymes (SBE) from maize (class A), pea (class A), maize (class B), rice (class B) and potato (class B), as well as human glycogen branching enzyme, allowed the inventors to identify a region in the carboxy-terminal one third of the protein which is almost completely conserved (GYLNFMGNEFGHPEWIDFPR) (Figure 4a). A multiple alignment of the DNA sequences (human, pea class A, potato class B, maize class B, maize class A and rice class B, respectively) corresponding to this region is shown in Figure 4b and was used to design an oligo which would potentially hybridize to all known plant starch branching enzymes: AATTT(C/T)ATGGGIAA(C/T)GA(A/G)TT(C/T)GG (Seq ID No. 20).

#### Library PCR

The initial isolation of a partial potato class A SBE cDNA clone was from an amplified potato tuber cDNA library in the  $\lambda$ Zap vector (Stratagene). One half  $\mu$ L of a potato cDNA library (titre  $2.3 \times 10^9$  pfu/mL) was used as template in a 50  $\mu$ L reaction containing 100 pmol of a 16 fold degenerate POTSBE primer and 25 pmol of a T7 primer (present in the  $\lambda$ Zap vector 3' to the cDNA sequences - see Figure 3), 100  $\mu$ M dNTPs, 2.5 U Taq polymerase and the buffer supplied with the Taq polymerase (Stratagene). All components except the enzyme were added to a 0.5 mL microcentrifuge tube, covered with mineral oil and incubated at 94°C for 7 minutes and then held at 55°C, while the Taq polymerase was added and mixed by pipetting. PCR was then performed by incubating for 1 min at 94°C, 1 min at 58°C and 3 minutes at 72°C, for 35 cycles. The PCR products were extracted with phenol/chloroform, ethanol precipitated and resuspended in TE pH 8.0 before cloning into the T/A cloning vector pT7BlueR (Invitrogen).

Several fragments between 600 and 1300 bp were amplified. These were isolated from an agarose gel and cloned into the pT7BlueR T/A cloning vector. Restriction mapping of 24 randomly selected clones showed that they belonged to several different groups (based on size and presence/absence of restriction sites). Initially four clones were chosen for sequencing. Of these four, two were found to correspond to the known potato class B SBE sequence, however the other two, although homologous, differed significantly and were more similar to the pea class A SBE sequence, suggesting that they belonged to the class A family of branching enzymes (Burton *et al.*, 1995 The Plant Journal, cited above). The latter two clones (~ 800bp) were sequenced fully. They both contained at the 5' end the sequence corresponding to the degenerate oligonucleotide used in the PCR and had a predicted open reading frame of 192 amino acids. The deduced amino acid sequence was highly homologous to that of the pea class A SBE.

The ~ 800 bp PCR derived cDNA fragment (corresponding to nucleotides 2281 to 3076 of the psbe2 con.seq sequence shown in Figure 8) was used as a probe to screen the potato tuber cDNA library. From one hundred and eighty thousand plaques, seven positives were obtained in the primary screen. PCR analysis showed that five of these clones were smaller than the original 800 bp cDNA clone, so these were not analysed further. The two other clones (designated 3.2.1 and 3.1.1) were approximately 1200 and 1500 bp in length respectively. These were sequenced from their 5' ends and the combined consensus sequence aligned with the sequence from the PCR generated clones. The cDNA clone 3.2.1 was excised from the phage vector and plasmid DNA was prepared and the insert fully sequenced. Several attempts to obtain longer clones from the library were unsuccessful, therefore clones containing the 5' end of the full length gene were obtained using RACE (rapid amplification of cDNA ends).

#### **Rapid Amplification of cDNA ends (RACE) and PCR conditions**

RACE was performed essentially according to Frohman (1992 Amplifications 11-15). Two  $\mu$ g of total RNA from mature potato tubers was heated to 65°C for 5 min and quick cooled on ice. The RNA was then reverse transcribed in a 20  $\mu$ L reaction for 1 hour at 37°C using BRL's M-MLV reverse transcriptase and buffer with 1 mM DTT, 1 mM dNTPs, 1 U/ $\mu$ L RNasin (Promega) and 500 pmol random hexamers (Pharmacia) as

primer. Excess primers were removed on a Centricon 100 column and cDNA was recovered and precipitated with isopropanol. cDNA was A-tailed in a volume of 20  $\mu$ l using 10 units terminal transferase (BRL), 200  $\mu$ M dATP for 10 min at 37°C, followed by 5 min at 65°C. The reaction was then diluted to 0.5 ml with TE pH 8 and stored at 4°C as the cDNA pool. cDNA clones were isolated by PCR amplification using the primers  $R_0R_1dT_{17}$ ,  $R_0$  and POTSBE24. The PCR was performed in 50  $\mu$ L using a hot start technique: 10  $\mu$ L of the cDNA pool was heated to 94°C in water for 5 min with 25 pmol POTSBE24, 25 pmol  $R_0$  and 2.5 pmol of  $R_0R_1dT_{17}$  and cooled to 75°C. Five  $\mu$ L of 10 x PCR buffer (Stratagene), 200  $\mu$ M dNTPs and 1.25 units of Taq polymerase were added, the mixture heated at 45°C for 2 min and 72°C for 40 min followed by 35 cycles of 94°C for 45 sec, 50°C for 25 sec, 72°C for 1.5 min and a final incubation at 72°C for 10 min. PCR products were separated by electrophoresis on 1% low melting agarose gels and the smear covering the range 600-800 bp fragments was excised and used in a second PCR amplification with 25 pmol of  $R_1$  and POTSBE25 primers in a 50  $\mu$ L reaction (28 cycles of 94°C for 1 min, 50°C 1 min, 72°C 2 min). Products were purified by chloroform extraction and cloned into pT7 Blue. PCR was used to screen the colonies and the longest clones were sequenced.

The first round of RACE only extended the length of the SBE sequence approximately 100 bases, therefore a new A-tailed cDNA library was constructed using the class A SBE specific oligo POTSBE24 (10 pmol) in an attempt to recover longer RACE products. The first and second round PCR reactions were performed using new class A SBE primers (POTSBE 28 and 29 respectively) derived from the new sequence data. Conditions were as before except that the elongation step in the first PCR was for 3 min and the second PCR consisted of 28 cycles at 94 °C for 45 seconds, 55 °C for 25 sec and 72 °C for 1 min 45 sec.

Clones ranging in size from 400 bp to 1.4 kb were isolated and sequenced. The combined sequence of the longest RACE products and cDNA clones predicted a full length gene of about 3150 nucleotides, excluding the poly(A) tail (psbe 2con.seq in Fig. 8).

As the sequence of the 5' half of the gene was compiled from the sequence of several

RACE products generated using Taq polymerase, it was possible that the compiled sequence did not represent that of a single mRNA species and/or had nucleotide sequence changes. The 5' 1600 bases of the gene was therefore re-isolated by PCR using Ultma, a thermostable DNA polymerase which, because it possesses a 3'-5' exonuclease activity, has a lower error rate compared to Taq polymerase. Several PCR products were cloned and restriction mapped and found to differ in the number of *Hind* III, *Ssp* I, and *Eco*R I sites. These differences do not represent PCR artefacts as they were observed in clones obtained from independent PCR reactions (data not shown) and indicate that there are several forms of the class A SBE gene transcribed in potato tubers.

In order to ensure that the sequence of the full length cDNA clone was derived from a single mRNA species it was therefore necessary to PCR the entire gene in one piece. cDNA was prepared according to the RACE protocol except that the adaptor oligo R<sub>0</sub>R<sub>1</sub>dT<sub>17</sub> (5 pmol) was used as a primer and after synthesis the reaction was diluted to 200  $\mu$ L with TE pH 8 and stored at 4°C. Two  $\mu$ L of the cDNA was used in a PCR reaction of 50  $\mu$ L using 25 pmol of class A SBE specific primers PBER1 and PBERT (see below), and thirty cycles of 94°C for 1 min, 60°C for 1 min and 72°C for 3 min. If Taq polymerase was used the PCR products were cloned into pT7Blue whereas if Ultma polymerase was used the PCR products were purified by chloroform extraction, ethanol precipitation and kinased in a volume of 20  $\mu$ L (and then cloned into pBSSK IIP which had been cut with *Eco*RV and dephosphorylated). At least four classes of cDNA were isolated, which again differed in the presence or absence of *Hind* III, *Ssp* I and *Eco*R I sites. Three of these clones were sequenced fully, however one clone could not be isolated in sufficient quantity to sequence.

The sequence of one of the clones (number 19) is shown in Figure 5. The first methionine (initiation) codon starts a short open reading frame (ORF) of 7 amino acids which is out of frame with the next predicted ORF of 882 amino acids which has a molecular mass (Mr) of approximately 100 Kd. Nucleotides 6-2996 correspond to SBE sequence - the rest of the sequence shown is vector derived. Figure 6 shows a comparison of the most highly conserved part of the amino acid sequence of potato class A SBE (residues 180-871, top, row) and potato class B SBE (bottom row, residues 98-792); the middle row indicates the

degree of similarity, identical residues being denoted by the common letter, conservative changes by two dots and neutral changes by a single dot. Dashes indicate gaps introduced to optimise the alignment. The class A SBE protein has 44% identity over the entire length with potato class B SBE, and 56% identity therewith in the central conserved domain (Figure 6), as judged by the "Megalign" program (DNASTAR). However, Figure 7 shows a comparison between potato class A SBE (top row, residues 1-873) and pea class A SBE (bottom row, residues 1-861), from which it can be observed that cloned potato gene is more homologous to the class A pea enzyme, where the identity is 70 % over nearly the entire length, and this increases to 83 % over the central conserved region (starting at IPPP at position -170). It is clear from this analysis that this cloned potato SBE gene belongs to the class A family of SBE genes.

An *E. coli* culture, containing the plasmid pSJ78 (which directs the expression of a full length potato SBE Class A gene), has been deposited (on 3rd January 1996) under the terms of the Budapest Treaty at The National Collections of Industrial and Marine Bacteria Limited (23 St Machar Drive, Aberdeen, AB2 1RY, United Kingdom), under accession number NCIMB 40781. Plasmid pSJ78 is equivalent to clone 19 described above. It represents a full length SBE A cDNA blunt-end ligated into the vector pBSSKIIIP.

#### Polymorphism of class A SBE genes

Sequence analysis of the other two full length class A SBE genes showed that they contain frameshift mutations and are therefore unable to encode full length proteins and indeed they were unable to complement the branching enzyme deficiency in the KV832 mutant (described below). An alignment of the full length DNA sequences is shown in Figure 8: "10con.seq" (Seq ID No. 12), "19con.seq" (Seq ID No. 14) and "11con.seq" (Seq ID No. 13) represent the sequence of full length clones 10, 19 and 11 obtained by PCR using the PBER1 and PBERT primers (see below), whilst "psbe2con.seq" (Seq ID No. 18) represents the consensus sequence of the RACE clones and cDNA clone 3.2.1. Those nucleotides which differ from the overall consensus sequence (not shown) are shaded. Dashes indicate gaps introduced to optimise the alignment. Apart from the frameshift mutations these clones are highly homologous. It should be noted that the 5' sequence of psbe2con is longer because this is the longest RACE product and it also contains several

changes compared to the other clones. The upstream methionine codon is still present in this clone but the upstream ORF is shortened to just 3 amino acids and in addition there is a 10 base deletion in the 5' untranslated leader.

The other significant area of variation is in the carboxy terminal region of the protein coding region. Closer examination of this area reveals a GAA trinucleotide repeat structure which varies in length between the four clones. These are typical characteristics of a microsatellite repeat region. The most divergent clone is #11 which has only one GAA triplet whereas clone 19 has eleven perfect repeats and the other two clones have five and seven GAA repeats. All of these deletions maintain the ORF but change the number of glutamic acid residues at the carboxy terminus of the protein.

Most of the other differences between the clones are single base changes. It is quite possible that some of these are PCR errors. To address this question direct sequencing of PCR fragments amplified from first strand cDNA was performed. Figure 9 shows the DNA sequence, and predicted amino acid sequence, obtained by such direct sequencing. Certain restriction sites are also marked. Nucleotides which could not be unambiguously assigned are indicated using standard IUPAC notation and, where this uncertainty affects the predicted amino acid sequence, a question mark is used. Sequence at the extreme 5' and 3' ends of the gene could not be determined because of the heterogeneity observed in the different cloned genes in these regions (see previous paragraph). However this can be taken as direct evidence that these differences are real and are not PCR or cloning artefacts.

There is absolutely no evidence for the frameshift mutations in the PCR derived sequence and it would appear that these mutations are an artefact of the cloning process, resulting from negative selection pressure in *E. coli*. This is supported by the fact that it proved extremely difficult to clone the full length PCR products intact as many large deletions were seen and the full length clones obtained were all cloned in one orientation (away from the LacZ promoter), perhaps suggesting that expression of the gene is toxic to the cells. Difficulties of this nature may have been responsible, at least in part, for the previous failure of other researchers to obtain the present invention.



A comparison of all the full length sequences is shown in Figure 10. In addition to clones 10, 11 and 19 are shown the sequences of a *Bgl* II - *Xho* I product cloned directly into the QE32 expression vector ("86CON.SEQ", Seq ID No. 16) and the consensus sequence of the directly sequenced PCR products ("pcrsbe2con.seq", Seq ID No. 17). Those nucleotides which differ from the consensus sequence (not shown) are shaded. Dashes indicate gaps introduced to optimise the alignment. There are 11 nucleotide differences predicted to be present in the mRNA population, which are indicated by asterisks above and below the sequence. The other differences are probably PCR artefacts or possibly sequencing errors.

#### Complementation of a branching enzyme deficient *E. coli* mutant

To determine if the isolated SBE gene encodes an active protein i.e. one that has branching enzyme activity, a complementation test was performed in the *E. coli* strain KV832. This strain is unable to make bacterial glycogen as the gene for the glycogen branching enzyme has been deleted (Keil *et al.*, 1987 Mol. Gen. Genet. 207, 294-301). When wild type cells are grown in the presence of glucose they synthesise glycogen (a highly branched glucose polymer) which stains a brown colour with iodine, whereas the KV832 cells make only a linear chain glucose polymer which stains blueish green with iodine. To determine if the cloned SBE gene could restore the ability of the KV832 cells to make a branched polymer, the clone pSJ90 (Seq ID No. 19) was used and constructed as below. The construct is a PCR-derived, substantially full length fragment (made using primers PBE 2B and PBE 2X, detailed below), which was cut with *Bgl* II and *Xho* I and cloned into the *Bam*H I / *Sal* I sites of the His-tag expression vector pQE32 (Qiagen). This clone, pSJ86, was sequenced and found to have a frameshift mutation of two bases in the 5' half of the gene. This frameshift was removed by digestion with *Nsi* I and *Sna*B I and replaced with the corresponding fragment from a Taq-generated PCR clone to produce the plasmid pSJ90 (sequence shown in Figure 12; the first 10 amino acids are derived from the expression vector). The polypeptide encoded by pSJ90 would be predicted to correspond to amino acids 46-882 of the full SBE coding sequence. The construct pSJ90 was transformed into the branching enzyme deficient KV832 cells and transformants were grown on solid PYG medium (0.85%  $\text{KH}_2\text{PO}_4$ , 1.1%  $\text{K}_2\text{HPO}_4$ , 0.6% yeast extract) containing 1.0% glucose. To test for complementation, a loop of cells was

scraped off and resuspended in 150 $\mu$ l of water, to which was added 15 $\mu$ l Lugol's solution (2g KI and 1g I<sub>2</sub> per 300ml water). It was found that the potato SBE fragment-transformed KV832 cells now stained a yellow-brown colour with iodine whereas control cells containing only the pQE32 vector continued to stain blue-green.

#### Expression of potato class A SBE in *E. coli*

Single colonies of KV832, containing one of the plasmids pQE32, pAGCR1 or pSJ90, were picked into 50ml of 2xYT medium containing carbenicillin, kanamycin and streptomycin as appropriate (100, 50 and 25 mg/L, respectively) in a 250ml flask and grown for 5 hours, with shaking, at 37°C. IPTG was then added to a final concentration of 1mM to induce expression and the flasks were further incubated overnight at 25°C. The cells were harvested by centrifugation and resuspended in 50 mM sodium phosphate buffer (pH 8.0), containing 300mM NaCl, 1mg/ml lysozyme and 1mM PMSF and left on ice for 1 hour. The cell lysates were then sonicated (3 pulses of 10 seconds at 40% power using a microprobe) and cleared by centrifugation at 12,000g for 10 minutes at 4°C. Cleared lysates were concentrated approximately 10 fold in a Centricon™ 30 filtration unit. Duplicate 10 $\mu$ l samples of the resulting extract were assayed for SBE activity by the phosphorylation stimulation method, as described in International Patent Application No. PCT/GB95/00634. In brief, the standard assay reaction mixture (0.2ml) was 200mM 2-(N-morpholino) ethanesulphonic acid (MES) buffer pH6.5, containing 100nCi of <sup>14</sup>C glucose-1-phosphate at 50mM, 0.05 mg rabbit phosphorylase A, and *E. coli* lysate. The reaction mixture was incubated for 60 minutes at 30°C and the reaction terminated and glucan polymer precipitated by the addition of 1ml of 75% (v/v) methanol, 1% (w/v) potassium hydroxide, and then 0.1ml glycogen (10mg/ml). The results are presented below:

Construct	SBE Activity (cpm)
pQE32 (control)	1,829
pSJ90 (potato class A SBE)	14,327
pAGCR1 (pea class A SBE)	29,707

The potato class A SBE activity is 7-8 fold above background levels. It was concluded therefore that the potato class A SBE gene was able to complement the BE mutation in the

phosphorylation stimulation assay and that the cloned gene does indeed code for a protein with branching enzyme activity.

### Oligonucleotides

The following synthetic oligonucleotides (Seq ID No.s 1-11 respectively) were used:

R <sub>0</sub> R <sub>1</sub> dT <sub>17</sub>	AAGGATCCGTCGACATCGATAATACGACTCACTATAGGGA(T) <sub>17</sub>
R <sub>0</sub>	AAGGATCCGTCGACATC
R <sub>1</sub>	GACATCGATAATACGAC
POTSBE24	CATCCAACCACCATCTCGCA
POTSBE25	TTGAGAGAAGATACCTAAGT
POTSBE28	ATGTTCAGTCCATCTAAAGT
POTSBE29	AGAACAACAATTCCTAGCTC
PBER 1	GGGGCCTTGAAGTCAGCAAT
PBERT	CGTCCCAGCATTCGACATAA
PBE 2B	CTTGGATCCTTGAAGTCAGCAATTTG
PBE 2X	TAAGTCGAGCAACGCGATCACAAGTTCGT

### Example 2

#### Production of Transgenic Plants

##### Construction of plant transformation vectors with antisense starch branching enzyme genes

A 1200 bp *Sac* I - *Xho* I fragment, encoding approximately the -COOH half of the potato class A SBE (isolated from the rescued  $\lambda$ Zap clone 3.2.1), was cloned into the *Sac* I - *Sal* I sites of the plant transformation vector pSJ29 to create plasmid pSJ64, which is illustrated schematically in Figure 11. In the figure, the black line represents the DNA sequence. The broken line represents the bacterial plasmid backbone (containing the origin of replication and bacterial selection marker), which is not shown in full. The filled triangles on the line denote the T-DNA borders (RB = right border, LB = left border). Relevant restriction sites are shown above the black line, with the approximate distances (in kilobases) between the sites (marked by an asterisk) given by the numerals below the

line. The thinnest arrows indicate polyadenylation signals (pAnos = nopaline synthase, pAg7 = Agrobacterium gene 7), the arrows intermediate in thickness denote protein coding regions (SBE II = potato class A SBE, HYG = hygromycin resistance gene) and the thickest arrows represent promoter regions (P-2x35 = double CaMV 35S promoter, Pnos = nopaline synthase promoter). Thus pSJ64 contained the class A SBE gene fragment in an antisense orientation between the 2X 35S CaMV promoter and the nopaline synthase polyadenylation signal.

For information, pSJ29 is a derivative of the binary vector pGPTV-HYG (Becker *et al.*, 1992 Plant Molecular Biology 20, 1195-1197) modified as follows: an approximately 750 bp (*Sac* I, T4 DNA polymerase blunted - *Sal* I) fragment of pJIT60 (Guerineau *et al.*, 1992 Plant Mol. Biol. 18, 815-818) containing the duplicated cauliflower mosaic virus (CaMV) 35S promoter (Cabb-JI strain, equivalent to nucleotides 7040 to 7376 duplicated upstream of 7040 to 7433, Frank *et al.*, 1980 Cell 21, 285-294) was cloned into the *Hind* III (Klenow polymerase repaired) - *Sal* I sites of pGPTV-HYG to create pSJ29.

#### Plant transformation

Transformation was conducted on two types of potato plant explants; either wild type untransformed minitubers (in order to give single transformants containing the class A antisense construct alone) or minitubers from three tissue culture lines (which gave rise to plants #12, #15, #17 and #18 indicated in Table 1) which had already been successfully transformed with the class B (SBE I) antisense construct containing the tandem 35S promoter (so as to obtain double transformant plants, containing antisense sequences for both the class A and class B enzymes).

Details of the method of Agrobacterium transformation, and of the growth of transformed plants, are described in International Patent Application No. WO 95/26407, except that the medium used contained 3% sucrose (not 1%) until the final transfer and that the initial incubation with Agrobacterium (strain 3850) was performed in darkness. Transformants containing the class A antisense sequence were selected by growth in medium containing 15mg/L hygromycin (the class A antisense construct comprising the HYG gene, i.e. hygromycin phosphotransferase).

Transformation was confirmed in all cases by production of a DNA fragment from the antisense gene after PCR in the presence of appropriate primers and a crude extract of genomic DNA from each regenerated shoot.

#### Characterisation of starch from potato plants

Starch was extracted from plants as follows: potato tubers were homogenised in water for 2 minutes in a Waring blender operating at high speed. The homogenate was washed and filtered (initially through 2mm, then through 1mm filters) using about 4 litres of water per 100gms of tubers (6 extractions). Washed starch granules were finally extracted with acetone and air dried.

Starch extracted from singly transformed potato plants (class A/SBE II antisense, or class B/SBE I antisense), or from double transformants (class A/SBE II and class B/SBE I antisense), or from untransformed control plants, was partially characterised. The results are shown in Table 1. The table shows the amount of SBE activity (units/gram tissue) in tubers from each transformed plant. The endotherm peak temperature (°C) of starch extracted from several plants was determined by DSC, and the onset temperature (°C) of pasting was determined by reference to a viscoamylograph ("RVA"), as described in WO 95/26407. The viscoamylograph profile was as follows: step 1 - 50°C for 2 minutes; step 2 - increase in temperature from 50°C to 95°C at a rate of 1.5°C per minute; step 3 - holding at 95°C for 15 minutes; step 4 - cooling from 95°C to 50°C at a rate of 1.5°C per minute; and finally, step 5 - holding at 50°C for 15 minutes. Table 1 shows the peak, pasting and set-back viscosities in stirring number units (SNU), which is a measure of the amount of torque required to stir the suspensions. Peak viscosity may be defined for present purposes as the maximum viscosity attained during the heating phase (step 2) or the holding phase (step 3) of the viscoamylograph. Pasting viscosity may be defined as the viscosity attained by the starch suspensions at the end of the holding phase (step 3) of the viscoamylograph. Set-back viscosity may be defined as the viscosity of the starch suspension at the end of step 5 of the viscoamylograph.

A determination of apparent amylose content (% w/w) was also performed, using the iodometric assay method of Morrison & Laignelet (1983 *J. Cereal Sci.* 1, 9-20). The

results (percentage apparent amylose) are shown in Table 1. The untransformed and transformed control plants gave rise to starches having apparent amylose contents in the range 29(+/-3)%.

Generally similar values for amylose content were obtained for starch extracted from most of the singly transformed plants containing the class A (SBE II) antisense sequence. However, some plants (#152, 249) gave rise to starch having an apparent amylose content of 37-38%, notably higher than the control value. Starch extracted from these plants had markedly elevated pasting onset temperatures, and starch from plant 152 also exhibited an elevated endotherm peak temperature (starch from plant 249 was not tested by DSC).

Table 1

Sample description	Sample number	Tuber SBE activity (µg starch)	DSC		Viscometry/ograph				Apparent amylose content (% w/w)	Phosphorus content (mg/100g)
			Peak temperature (°C)	Onset temperature (°C)	Peak viscosity (cStU)	Pasting viscosity (cStU)	Set-back viscosity (cStU)			
Untransformed control	146	7.8	65.8	65.8	645	161	200	31.2	66	
	243	22.2	nd	62.8	761	135	241	20.1		
AS-Class A SBE	152	12.7	68.5	70.8	467	380	528	37.5	88	
	248	13.8	nd	70.0	487	434	616	38.5		
AS-Class B SBE (17) (control)	145	0.7	68.8	68.8	688	177	305	28.8	111	
AS-Class B SBE (17) + AS-Class A SBE	150	0.6	74.0	68.0	214	214	303	53.1	188	
	161	0.5	73.0	78.8	348	324	616	40.8	208	
AS-Class B SBE (18) (control)	144	1.8	64.5	64.7	714	154	258	26.0	87	
AS-Class B SBE (18) + AS-Class A SBE	149	3.0	68.5	68.8	474	287	482	35.8	127	
AS-Class B SBE (18) (control)	172	0.22	nd	63.4	707	187	280	28.8	130	
AS-Class B SBE (18) + AS-Class A SBE	201	0.10	nd	>85	no peak	12	13	88.4	210	
	206a	0.10	nd	>85	no peak	15	17	84.1		
	208	0.30	72.8-80.5	>85	no peak	14	19	82.8	240	
	202	0.02	nd	88.4	no peak	172	245	67.8		
	212	1.40	nd	78.0	308	288	541	48.5		
	220	1.40	nd	73.8	355	345	583	44.1		
AS-Class B SBE (12) (control)	170	0.2	nd	68.5	768	202	303	27.8		
AS-Class B SBE (12) + AS-Class A SBE	236	0.7	nd	85.0	no peak	23	14	80.4		
	236a	0.9	nd	91.2	no peak	138	182	58.7		
	230a	0.8	nd	77.6	244	238	450	48.2		

RVA profile

Pasting viscosity (47 min)

Set back viscosity (92 min)

SBE

SNU

nd

50°C (2 min), 50-60°C (1.5°C/min), 65°C (15 min), 65-80°C (1.5°C/min), 80°C (18 min)

at end of 50°C (2 min), 50-60°C (1.5°C/min), 65°C (15 min)

at end of profile

Starch Branching Enzyme

Instrument "Starch Number Unit" (arbitrary units)

not determined

Table 1

Sample description	Sample number	Tuber SBE activity (U/g starch)	DSC	
			Peak temperature (°C)	Onset temperature (°C)
Untransformed control	146	7.6	65.8	65.5
	243	22.2	nd	62.6
AS-Class A SBE	152	12.7	68.5	70.9
	249	13.9	nd	70.0
AS-Class B SBE (17) (control)	145	0.7	68.9	66.8
AS-Class B SBE (17) + AS-Class A SBE	150	0.6	74.0	86.0
	161	0.5	73.0	76.6
AS-Class B SBE (18) (control)	144	1.6	64.5	64.7
AS-Class B SBE (18) + AS-Class A SBE	149	3.0	68.5	69.9

SUBSTITUTE SHEET (RULE 26)



29/2

Viscoamylograph			(RVA)		Apparent amylose content (% w/w)	Phosphorus content (mg/100g)
Peak viscosity (SNU)	Pasting viscosity (SNU)	Set-back viscosity (SNU)				
545	161	280	68	31.2	68	68
761	135	241				
467	380	529	89	37.5	89	89
497	434	518				
669	177	305	111	29.8	111	111
214	214	303				
349	324	618	198	53.1	198	198
714	154	258				
474	267	482	206	40.9	206	206
			97	29.0	97	97
			127	35.6	127	127

SUBSTITUTE SHEET (RULE 26)

AS-Class B SBE (15) (control)	172	0.22	nd	65.4
AS-Class B SBE (15) + AS-Class A SBE	201	0.10	nd	>95
	208a	0.10	nd	>95
	208	0.30	72.8-80.5	>95
	202	0.02	nd	89.4
	212	1.40	nd	78.0
	220	1.40	nd	75.8
AS-Class B SBE (12) (control)	170	0.2	nd	86.5
AS-Class B SBE (12) + AS-Class A SBE	238	0.7	nd	95.0
	238a	0.9	nd	91.2
	230a	0.8	nd	77.6

RVA profile

Pasting viscosity (47 min)

Set-back viscosity (92 min)

SBE

SNU

nd

50°C (2 min), 50-95°C (1.5°C/min), 95°C (15 min), 95-50°C (1.5°C/min), 50°C (15 min)

at end of 50°C (2min), 50-95°C (1.5°C/min), 95°C (15 min)

at end of profile

Starch Branching Enzyme

Instrument "Stirring Number Units" (arbitrary units)

not determined

SUBSTITUTE SHEET (RULE 26)

29/4

707	167	280	28.8	130
no peak	12	13	66.4	210
no peak	15	17	64.1	
no peak	14	19	62.8	240
no peak	172	245	57.9	
308	296	541	49.5	
355	345	593	44.1	
768	202	303	27.8	
no peak	23	14	60.4	
no peak	139	192	56.7	
244	239	450	48.2	

SUBSTITUTE SHEET (RULE 26)

It should be noted that, even if other single transformants were not to provide starch with an altered amylose/amylopectin ratio, the starch from such plants might still have different properties relative to starch from conventional plants (e.g. different average molecular weight or different amylopectin branching patterns), which might be useful.

Double transformant plants, containing antisense sequences for both the class A and class B enzymes, had greatly reduced SBE activity (units/gm) compared to untransformed plants or single anti-sense class A transformants, (as shown in Table 1). Moreover, certain of the double transformant plants contained starch having very significantly altered properties. For example, starch extracted from plants #201, 202, 208, 208a, 236 and 236a had drastically altered amylose/amylopectin ratios, to the extent that amylose was the main constituent of starch from these plants. The pasting onset temperatures of starch from these plants were also the most greatly increased (by about 25-30°C). Starch from plants such as #150, 161, 212, 220 and 230a represented a range of intermediates, in that such starch displayed a more modest rise in both amylose content and pasting onset temperature. The results would tend to suggest that there is generally a correlation between % amylose content and pasting onset temperature, which is in agreement with the known behaviour of starches from other sources, notably maize.

The marked increase in amylose content obtained by inhibition of class A SBE alone, compared to inhibition of class B SBE alone (see PCT/GB95/00634) might suggest that it would be advantageous to transform plants first with a construct to suppress class A SBE expression (probably, in practice, an antisense construct), select those plants giving rise to starch with the most altered properties, and then to re-transform with a construct to suppress class B SBE expression (again, in practice, probably an antisense construct), so as to maximise the degree of starch modification.

In addition to pasting onset temperatures, other features of the viscoamylograph profile e.g. for starches from plants #149, 150, 152, 161, 201, 236 and 236a showed significant differences to starches from control plants, as illustrated in Figure 13. Referring to Figure 13, a number of viscoamylograph traces are shown. The legend is as follows: shaded box - normal potato starch control (29.8% amylose content); shaded circle - starch from plant

149 (35.6% amylose): shaded triangle, pointing upwards - plant 152 (37.5%); shaded triangle, pointing downwards - plant 161 (40.9%); shaded diamond - plant 150 (53.1%); unshaded box - plant 236a (56.7%); unshaded circle - plant 236 (60.4%); unshaded triangle, pointing upwards - plant 201 (66.4%); unshaded triangle, pointing downwards - Hylon V starch, from maize (44.9 % amylose). The thin line denotes the heating profile.

With increasing amylose content, peak viscosities during processing to 95°C decrease, and the drop in viscosity from the peak until the end of the holding period at 95°C also generally decreases (indeed, for some of the starch samples there is an increase in viscosity during this period). Both of these results are indicative of reduced granule fragmentation, and hence *increased* granule stability during pasting. This property has not previously been available in potato starch without extensive prior chemical or physical modification. For applications where a maximal viscosity after processing to 95°C is desirable (i.e. corresponding to the viscosity after 47 minutes in the viscoamylograph test), starch from plant #152 would be selected as starches with both lower (Controls, #149) and higher (#161, #150) amylose contents have lower viscosities following this gelatinisation and pasting regime (Figure 13 and Table 1). It is believed that the viscosity at this stage is determined by a combination of the extent of granule swelling and the resistance of swollen granules to mechanical fragmentation. For any desired viscosity behaviour, one skilled in the art would select a potato starch from a range containing different amylose contents produced according to the invention by performing suitable standard viscosity tests.

Upon cooling pastes from 95°C to 50°C, potato starches from most plants transformed in accordance with the invention showed an increase in viscoamylograph viscosity as expected for partial reassociation of amylose. Starches from plants #149, 152 and 161 all show viscosities at 50°C significantly in excess of those for starches from control plants (Figure 13 and Table 1). This contrasts with the effect of elevated amylose contents in starches from maize plants (Figure 2) which show very low viscosities throughout the viscoamylograph test. Of particular note is the fact that, for similar amylose contents, starch from potato plant 150 (53 % amylose) shows markedly increased viscosity compared with Hylon 5 starch (44.9% amylose) as illustrated in Figure 13. This demonstrates that

useful properties which require elevated (35% or greater) amylose levels can be obtained by processing starches from potato plants below 100°C, whereas more energy-intensive processing is required in order to generate similarly useful properties from high amylose starches derived from maize plants.

Final viscosity in the viscoamylograph test (set-back viscosity after 92 minutes) is greatest for starch from plant #161 (40.9% amylose) amongst those tested (Figure 13 and Table 1). Decreasing final viscosities are obtained for starches from plant #152 (37.5% amylose), #149 (35.6% amylose) and #150 (53.1% amylose). Set-back viscosity occurs where amylose molecules, exuded from the starch granule during pasting, start to re-associate outside the granule and form a viscous gel-like substance. It is believed that the set-back viscosity values of starches from transgenic potato plants represent a balance between the inherent amylose content of the starches and the ability of the amylose fraction to be exuded from the granule during pasting and therefore be available for the reassociation process which results in viscosity increase. For starches with low amylose content, increasing the amylose content tends to make more amylose available for re-association, thus increasing the set-back viscosity. However, above a threshold value, increased amylose content is thought to inhibit granule swelling, thus preventing exudation of amylose from the starch granule and reducing the amount of amylose available for re-association. This is supported by the RVA results obtained for the very high amylose content potato starches seen in the viscoamylograph profiles in Figure 13. For any desired viscosity behaviour following set-back or retrogradation to any desired temperature over any desired timescale, one skilled in the art would select a potato starch from a range containing different amylose contents produced according to the invention by performing standard viscosity tests.

Further experiments with starch from plants #201 and 208 showed that this had an apparent amylose content of over 62% (see Table 1). Viscoamylograph studies showed that starch from these plants had radically altered properties and behaved in a manner similar to hylon 5 starch from maize plants (Figure 13). Under the conditions employed in the viscoamylograph, this starch exhibited extremely limited (nearly undetectable) granule swelling. Thus, for example, unlike starch from control plants, starch from plants

201, 208 and 208a did not display a clearly defined pasting viscosity peak during the heating phase. Microscopic analysis confirmed that the starch granule structure underwent only minor swelling during the experimental heating process. This property may well be particularly useful in certain applications, as will be apparent to those skilled in the art.

Some re-grown plants have so far been found to increase still further the apparent amylose content of starch extracted therefrom. Such increases may be due to:-

- i) Growth and development of the first generation transformed plants may have been affected to some degree by the exogenous growth hormones present in the tissue culture system, which exogenous hormones were not present during growth of the second generation plants; and
- ii) Subsequent generations were grown under field conditions, which may allow for attainment of greater maturity than growth under laboratory conditions, it being generally held that amylose content of potato starch increases with maturity of the potato tuber.

Accordingly, it should be possible to obtain potato plants giving rise to tubers with starch having an amylose content in excess of the 66% level so far attained, simply by analysing a greater number of transformed plants and/or by re-growing transgenic plants through one or more generations under field conditions.

Table 1 shows that another characteristic of starch which is affected by the presence of anti-sense sequences to SBE is the phosphorus content. Starch from untransformed control plants had a phosphorus content of about 60-70mg/100gram dry weight (as determined according to the AOAC Official Methods of Analysis, 15th Edition, Method 948.09 "Phosphorus in Flour"). Introduction into the plant of an anti-sense SBE B sequence was found to cause a modest increase (about two-fold) in phosphorus content, which is in agreement with the previous findings reported at scientific meetings. Similarly, anti-sense to SBE A alone causes only a small rise in phosphorus content relative to untransformed controls. However, use of anti-sense to both SBE A and B in combination results in up to a four-fold increase in phosphorus content, which is far greater than any *in planta* phosphorus content previously demonstrated for potato starch.

This is useful in that, for certain applications, starch must be phosphorylated *in vitro* by

chemical modification. The ability to obtain potato starch which, as extracted from the plant, already has a high phosphorus content will reduce the amount of *in vitro* phosphorylation required suitably to modify the starch. Thus, in another aspect the invention provides potato starch which, as extracted from the plant, has a phosphorus content in excess of 200mg/100gram dry weight starch. Typically the starch will have a phosphorus content in the range 200 - 240mg/100gram dry weight starch.



## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: National Starch and Chemical Investment Holding Corporation
- (B) STREET: 501 Silverside Road, Suite 27
- (C) CITY: Wilmington
- (D) STATE: Delaware
- (E) COUNTRY: United States of America
- (F) POSTAL CODE (ZIP): 19809

(ii) TITLE OF INVENTION: Improvements in or Relating to Plant Starch Composition

(iii) NUMBER OF SEQUENCES: 20

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0. Version #1.30 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

AAGGATCCGT CGACATCGAT AATACGACTC ACTATAGGGA TTTTTTTTTT TTTTTT

57

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

AAGGATCCGT CGACATC

17

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs

36

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GACATCGATA ATACGAC

17

(2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CATCCAACCA CCATCTCGCA

20

(2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

TTGAGAGAAG ATACCTAAGT

20

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

ATGTTCAAGTC CATCTAAAGT

20

(2) INFORMATION FOR SEQ ID NO: 7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

AGAACAACAA TTCCTAGCTC

20

(2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GGGGCCTTGA ACTCAGCAAT

20

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

CGTCCCAGCA TTCGACATAA

20

(2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

CTTGGATCCT TGA ACTCAGC AATTTG

26

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

TAACTCGAGC AACGCGATCA CAAGTTCGT

29

## (2) INFORMATION FOR SEQ ID NO: 12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3003 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

GATGGGGCCT TGAACTCAGC AATTGACAC TCAGTTAGTT ACACTGCCAT CACTTATCAG	60
ATCTCTATTT TTTCTCTTAA TTCCAACCAA GGAATGAATA AAAAGATAGA TTTGTAAAAA	120
CCCTAAGGAG AGAAGAAGAA AGATGGTGTA TAACTCTCT GGAGTTCGTT TTCCTACTGT	180
TCCATCAGTG TACAAATCTA ATGGATTGAG CAGTAATGGT GATCGGAGGA ATGCTAATAT	240
TTCTGTATTC TTGAAAAAAC ACTCTCTTTC ACGGAAGATC TTGGCTGAAA AGTCTTCTTA	300
CAATTCCGAA TCCCGACCTT CTACAATTGC AGCATCGGGG AAAGTCCTTG TGCCTGGAAT	360
CCAGAGTGAT AGCTCCTCAT CCTCAACAGA TCAATTTGAG TTCGCTGAGA CATCTCCAGA	420
AAATCCCCCA GCATCAACTG ATGTAGATAG TTCAACAATG GAACACGCTA GCCAGATTAA	480
AACTGAGAAC GATGACGTTG AGCCGTCAAG TGATCTTACA GGAAGTGTTG AAGAGCTGGA	540
TTTTGCTTCA TCACTACAAC TACAAGAAGG TGGTAACTG GAGGAGTCTA AAACATTAAA	600
TACTTCTGAA GAGACAATTA TTGATGAATC TGATAGGATC AGAGAGAGGG GCATCCCTCC	660
ACCTGGACTT GGTGAGAAGA TTTATGAAAT AGACCCCTT TTGACAACT ATCGTCAACA	720
CCTTGATTAC AGGTATTCAC AGTACAAGAA ACTGAGGGAG GCAATTGACA AGTATGAGGG	780
TGGTTTGGAA GCTTTTTCTC GTGGTTATGA AAGAATGGGT TTCACTCGTA GTGCTACAGG	840
TATCACTTAC CGTGAGTGGG CTCCTGGTGC CCAGTCAGCT GCCCTCATTG GGGATTTCAA	900
CAATTGGGAC GCAAATGCTG ACTTTATGAC TCGGAATGAA TTTGGTGTCT GAGAGATTTT	960
TCTGCCAAAT AATGTGGATG GTTCTCCTGC AATTCCTCAT GGGTCCAGAG TGAAGATACG	1020
TATGGAACT CCATCAGGTG TTAAGGATTC CATTCTGCT TGGATCACT ACTCTTTACA	1080
GCTTCCTGAT GAAATTCCAT ATAATGGAAT ATATTATGAT CCACCCGAAG AGGAGAGGTA	1140
TATCTTCAA CACCCACGGC CAAAGAAACC AAAGTCGGTG AGAATATATG AATCTCATAT	1200
TGGAATGAGT AGTCCGGAGC CTAAATTAA CTCATACGTG AATTTTAGAG ATGAAGTTCT	1260
TCCTGCATA AAAAAAGCTT GGGTACAATG CGGTGCAAAT TATGGCTATT CAAGAGCATT	1320
CTTATTATGC TAGTTTTGGT TATCATGTCA CAAATTTTTT TGCACCAAGC AGCCGTTTTG	1380

GAACGCCCCGA CGACCTTAAG TCTTTGATTG ATAAAGCTCA TGAGCTAGGA ATTGTTGTTC	1440
TCATGGACAT TGTTACAGC CATGCATCAA ATAATACTTT AGATGGACTG AACATGTTTG	1500
ACGGCACAGA TAGTTGTTAC TTTCACTCTG GAGCTCGTGG TTATCATTGG ATGTGGGATT	1560
TCCGCTCTT TAACTATGGA AACTGGGAGG TACTTAGGTA TCTTCTCTCA AATGCGAGAT	1620
GGTGGTTGGA TGAGTTCAAA TTTGATGGAT TTAGATTGA TGGTGTGACA TCAATGATGT	1680
GTA CTCACCA CGGATTATCG GTGGGATTCA CTGGGA ACTA CGAGGAATAC TTTGGACTCG	1740
CAACTGATGT GGATGCTGTT GTGTATCTGA TGCTGGTCAA CGATCTTATT CATGGGCTTT	1800
TCCCAGATGC AATTACCATT GGTGAAGATG TTAGCGGAAT GCCGACATTT TGTGTTCCCG	1860
TTCAAGATGG GGGTGTGGC TTTGACTATC GGCTGCATAT GGCAATTGCT GATAAATGGA	1920
TTGAGTTGCT CAAGAAACGG GATGAGGATT GGAGAGTGGG TGATATTGTT CATACACTGA	1980
CAAATAGAAG ATGGTCGGAA AAGTGTGTTT CATACGCTGA AAGTCATGAT CAAGCTCTAG	2040
TCCGTGATAA AACTATAGCA TTCTGGCTGA TGGACAAGGA TATGTATGAT TTTATGGCTC	2100
TGGATAGACC GTCAACATCA TTAATAGATC GTGGGATAGC ATTACACAAG ATGATTAGGC	2160
TTGTA ACTAT GGGATTAGGA GGAGAAGGGT ACCTAAATTT CATGGGAAAT GAATTCGGCC	2220
ACCCTGAGTG GATTGATTTT CCTAGGGCTG AACACACCT CTCTGATGGC TCAGTAATTC	2280
CCAGAAACCA ATTCAGTTAT GATAAATGCA GACGGAGATT TGACCTGGGA GATGCAGAAT	2340
ATTTAAGATA CCGTGGGTTG CAAGAATTTG ACCGGGCTAT GCAGTATCTT GAAGATAAAT	2400
ATGAGTTTAT GACTTCAGAA CACCAGTTCA TATCACGAAA GGATGAAGGA GATAGGATGA	2460
TTGTATTTGA AAAAGGAAAC CTAGTTTTTG TCTTTAATTT TCACTGGACA AAAGGCTATT	2520
CAGACTATCG CATAGGCTGC CTGAAGCCTG GAAAATACAA GGTTCCTTG GACTCAGATG	2580
ATCCACTTTT TGGTGGCTTC GGGAGAATTG ATCATAATGC CGAATATTTT ACCTTTGAAG	2640
GATGGTATGA TGATCGTCCT CGTTCAATTA TGGTGTATGC ACCTAGTAGA ACAGCAGTGG	2700
TCTATGCACT AGTAGACAAA GAAGAAGAAG AAGAAGAAGA AGTAGCAGTA GTAGAAGAAG	2760
TAGTAGTAGA AGAAGAATGA ACGAACTTGT GATCGCGTTG AAAGATTTGA ACGCCACATA	2820
GAGCTTCTTG ACGTATCTGG CAATATTGCA TTAGTCTTGG CGGAATTTCA TGTGACAACA	2880
GGTTTGCAAT TCTTTCCACT ATTAGTAGTG CAACGATATA CGCAGAGATG AAGTGCTGAA	2940
CAAAAACATA TGTA AAATCG ATGAATTTAT GTCGAATGCT GGGACGATCG AATTCCTGCA	3000
GCC	3003

20/75

↙180 ↙190 ↙200 ↙210 ↙220  
 IYEIDPLLTNYRQHLDYRYSQYKKLREAIIDKYEGGLEAFSRGYEKMGR  
 : : DP L. Y : H: . R : : Y : : I: KYEG LE. F: : GY K. GF. R  
 LLNLDPTLEPYLDHFRHRMKRYVDQKMLIEKYEGPLEEFAQGYLKFGFNR  
 ↗100 ↗110 ↗120 ↗130 ↗140  
 ↙230 ↙240 ↙250 ↙260 ↙270  
 SATGITYREWALGAQSAALIGDFNNWDANADIMTRNEFGVWEIFLPNNVD  
 ... I. YREWA : AQ. A. : IGDFN. W: : : : M. : : : FGVW. I : P: VD  
 EDGCIVYREWAPAAQEAIEVIGDFNGWNGSNHMEKDQFGVWSIRIPD-VD  
 ↗150 ↗160 ↗170 ↗180 ↗190  
 ↙280 ↙290 ↙300 ↙310 ↙320  
 GSPAIPHGSRVKIRMDTPSGV-KDSIPAWINYSLQLPDEI--PYNGIHYD  
 : . P. IPH. SRVK: R. : : GV D. IPAWI: Y: : : : PY: G: . D  
 SKPVIPHNSRVKFRFKHNGVWVDRIPAWIKYATADATKFAAPYDGVYWD  
 ↗200 ↗210 ↗220 ↗230 ↗240  
 ↙330 ↙340 ↙350 ↙360 ↙370  
 PPEEERYIFQHPRPKPKSLRIYESHIGMSSPEPKINSYVNRDEVLPR  
 PP . ERY F: . PRP KP: : RIYE: H: GMSS: EP: : NSY : F D: VLPRI  
 PPPSERYHFKYPRPPKPRAPRIYEAHVGMSSSEPRVNSYREFADDVLPR  
 ↗250 ↗260 ↗270 ↗280 ↗290  
 ↙380 ↙390 ↙400 ↙410 ↙420  
 KKLGYNALQIMAIQEHSSYYASFGYHVTNFFAPSSRFGTPDDLKSLIDKAH  
 K . YN: : Q: MAI EHSYY: SFGYHVTNFFA S: R: G: P: DLK LIDKAH  
 KANNYNTVOLMAIMEHSYYGSFGYHVTNFFAVSNRYGNPEDLKYLIDKAH  
 ↗300 ↗310 ↗320 ↗330 ↗340  
 ↙430 ↙440 ↙450 ↙460 ↙470  
 ELGIVVLMDIVHSHASNNTLDGLNMFDC---TDSCYFHSGARGYHWMWDS  
 . LG: VL: D: VHSHASN. DGLN FD : : : YFH: G. RGYH : WDS  
 SLGLQVLVDVVHSHASNNTDGLNGFDIGOGSDESYFHAGERGYHKLWDS  
 ↗350 ↗360 ↗370 ↗380 ↗390  
 ↙480 ↙490 ↙500 ↙510 ↙520  
 RLFNYGNWEVLRLLSNARWWLDAFKFDGFRFDGVTSMYIHHGLSVGFT  
 RLFNY: NWEVLR: LLSN RWL: : : FDGFRFDG: TSM: Y: HHG: : : GFT  
 RLFNYANWEVLRFLLSNLRWWLEENFDGFRFDGITSMLYVHHGINMGFT  
 ↗400 ↗410 ↗420 ↗430 ↗440  
 ↙530 ↙540 ↙550 ↙560 ↙570  
 GNYEEYFGLATDVDVAVVYLMVNDLIHGLFPDAITIGEDVSGMPTFCIPV  
 GNY: EYF: ATDVDVAVVYLM. N: LIH : FPDA. . I: EDVSGMP. : . PV  
 GNYNEYFSEATDVDVAVVYLMANLIHKIFPDATVIAEDVSGMPGLSRPV  
 ↗450 ↗460 ↗470 ↗480 ↗490  
 ↙580 ↙590 ↙600 ↙610 ↙620  
 QEGGVGFDYRLHMAIADKRIELLK-KRDEDWRVGDIVHTLTNRRWSEKCV  
 EGG: GFDYRL MAI: DK: I: LK K. DEDW: : : : LTNRR: : EKC:  
 SEGGIGFDYRLAMAIIPDKWIDYLNKNDEDSMKEVTSSLTNRRYTEKCI  
 ↗500 ↗510 ↗520 ↗530 ↗540

Fig. 6 SHEET 1

SUBSTITUTE SHEET (RULE 26)

21/75

✓630      ✓640      ✓650      ✓660      ✓670  
 SYAESHDOALVGDKTIAFWLMDKDMYDFMALDRPSTSLIDRGIALHKMIR  
 : YAESHDOQ: VGDKTIAF LMDK: MY. M: ::::: DRGIALHKMI:  
 AYAESHDOQIVGDKTIAFLMDKEMYSGMSCLTDASPVVDRGIALHKMIH  
 ^550      ^560      ^570      ^580      ^590  
 ✓680      ✓690      ✓700      ✓710      ✓720  
 LVTMGLGGEGYLNFMGNEFGHPEWIDFPRAEQHLSDGSVIPGNQFSYDKC  
 : TM: LGGEGYLNFMGNEFGHPEWIDFPR      GN: SYDKC  
 FFTMALGGEGYLNFMGNEFGHPEWIDFPR-----EGNNWSYDKC  
 ^600      ^610      ^620      ^630  
 ✓730      ✓740      ✓750      ✓760      ✓770  
 RRRFDLGDAEYLR YRGLQEFDRPMQYLEDKYEFMTSEHQFISRKDEGDRM  
 RR: : L: D: E. LRY: : : FDR: M: L: : K: : F: : S: : Q: : S: : D: : : :  
 RROWNLA DSEHLRYKFMNAFDRAMNSLDEKFSFLASGKQIVSSMDDDNKV  
 ^640      ^650      ^660      ^670      ^680  
 ✓780      ✓790      ✓800      ✓810      ✓820  
 IVFEKGNLVFVFNFWTKSYSDYRIACLKPGKYKVALDSDDPLFGGFGRI  
 : VFE: G: LVFVFNFH : : Y: : Y: : : C PGKY: VAL: SD. FGG GR  
 VVFERGDLVFVFNFWHPNNTYEGYKVGCDLPGKYRVALGSDAWEFGGHGRA  
 ^690      ^700      ^710      ^720      ^730  
 ✓830      ✓840      ✓850      ✓860  
 DHNAEYFT-----FEGWYDDRPRSIMVYAPCKTAVVYALVDKEEEEE  
 : H: : : FT      E. : : : RP. S: : V : P : T V. Y VD. : E.  
 GHDVDHFTSPEGIPGVPETNFNGRPNSEFKVLSPARTCVAYYRVDERMSET  
 ^740      ^750      ^760      ^770      ^780  
 ✓870  
 EEEEEEV  
 E: : : :  
 EDYQTDI  
 ^790

Fig. 6 SHEET 2

SUBSTITUTE SHEET (RULE 26)

22/75

M V Y T L S G V R F P T V P S V Y K S N G F S S N G D R R N A N V S V F L K K H -- S L S R K I L A  
 M V Y T : S G : R F P : : P S : . K S : . . D R R : : : S F L K : : : S : S R . L  
 M V Y T I S G I R F P V L P S L H K S --- T L R C D R R A S S H S F F L K N N S S S F S R T S L Y  
 E K S S Y N S E F R P S T V A A S G K V L V P G T Q S D S S S S S T D Q F E F T E T S P E N S P A S  
 . K S : S E : : S T : A . S : K V L : P . . Q D : S S : D Q : E . . . . : E : . . .  
 A K F S R D S E T K S S T I A E S D K V L I P E D Q - D N S V S L A D Q L E N P D I T S E D A Q N L  
 T D V D S S T M E H A S Q I K T E N D D V E P S S D L T G S V E E L D F A S S L O L Q E G G K L E E  
 . D : T M . : : : : : : : : : : : : : : : : S : : : : : : :  
 E D L --- T M K D G N K Y N I D - E S T S S Y R E V G D E K G S V T S S S L V D V N T D T Q -- A  
 S K T L N T S E E T I I D E S D R I R E R G I P P P G L G Q K I Y E I D P L L T N Y R O H L D Y R Y  
 . K T S : . . . : . . : I I P P P G G Q K I Y E I D P L L . . R O H L D : R Y  
 K K T S V H S D K K V K V D K P K I --- I P P P G S G Q K I Y E I D P L L Q A H R O H L D F R Y  
 S O Y K K L R E A I D K Y E G G L E A F S R G Y E K M G F T R S A T G I T Y R E W A L G A Q S A A L  
 : O Y K : : R E . I D K Y E G G L : A F S R G Y E K . G F T R S A T G I T Y R E W : G A : S A A L  
 G O Y K R I R E E I D K Y E G G L D A F S R G Y E K F G F T R S A T G I T Y R E W G P G A K S A A L  
 I G D F N N W D A N A D I M T R N E F G V W E I F L P N N V D G S P A I P H G S R V K I R M D T P S  
 : G D F N N W : : N A D : M T : : . F G V W E I F L P N N . D G S P : I P H G S R V K I : M D T P S  
 V G D F N N W N P N A D V M T K D A F G V W E I F L P N N A D G S P P I P H G S R V K I H M D T P S  
 G V K D S I P A W I N Y S L Q L P D E I P Y N G I H Y D P P E E E R Y I F O H P R P K K P K S L R I  
 G : K D S I P A W I : : S : Q P : E I P Y N G I . Y D P P E E E : Y : F : H P : P K : P : S : R I  
 G I K D S I P A W I K F S V Q A P G E I P Y N G I Y Y D P P E E E K Y V F K H P O P K R P Q S I R I  
 Y E S H I G M S S P E P K I N S Y V N F R D E V L P R I K K L G Y N A L Q I M A I Q E H S Y Y A S F  
 Y E S H I G M S S P E P K I N : Y . N F R D : V L P R I K K L G Y N A : Q I M A I Q E H S Y Y A S F  
 Y E S H I G M S S P E P K I N T Y A N F R D D V L P R I K K L G Y N A V Q I M A I Q E H S Y Y A S F  
 G Y H V T N F F A P S S R F G T P D D L K S L I D K A H E L G I V V L M D I V H S H A S N N T L D G  
 G Y H V T N F F A P S S R F G T P : D L K S L I D : A H E L G : : V L M D I V H S H : S N N T L D G  
 G Y H V T N F F A P S S R F G T P E D L K S L I D R A H E L G L L V L M D I V H S H S S N N T L D G

Fig. 7 SHEET 1

SUBSTITUTE SHEET (RULE 26)



23/75

↙450 ↙460 ↙470 ↙480 ↙490  
 LNMFDCTDSCYFHSGARGYHWMWDSRLFNYSWEVRLRYLLSNARWWLDAF  
 LNMFD TD: YFH: G: RGYHWMWDSRLFNYSWEVRLRYLLSNARWWLD. :  
 LNMFDGTDGHYFHPGSRGYHWMWDSRLFNYSWEVRLRYLLSNARWWLDEY  
 ↗440 ↗450 ↗460 ↗470 ↗480  
 ↙500 ↙510 ↙520 ↙530 ↙540  
 KFDGFRFDGVTSMYIHHGLSVGFTGNYEEYFGLATDVEDAVVYMLVNDL  
 KFDGFRFDGVTSMY. HHGL V: FTGNY. EYFGLATDV: AVVY: MLVNDL  
 KFDGFRFDGVTSMYTHHGLQVSFTGNYSEYFGLATDVEAVVYMLVNDL  
 ↗490 ↗500 ↗510 ↗520 ↗530  
 ↙550 ↙560 ↙570 ↙580 ↙590  
 IHGLFPDAITIGEDVSGMPTFCIPVQEGGVGFDYRLHMAIADKRIELLKK  
 IHGLFP: A: : IGEDVSGMPTFC: P. Q: GG: GF: YRLHMA: ADK: IELLKK  
 IHGLFPEAVSIGEDVSGMPTFCLPTQDGGIGFNRYRLHMAVADKWIELLKK  
 ↗540 ↗550 ↗560 ↗570 ↗580  
 ↙600 ↙610 ↙620 ↙630 ↙640  
 RDEDWRVGDIVHTLTNRRWSEKCVSYAESHDQALVGDKTIAFWLMDKDMY  
 : DEDWR: GDIVHTLTNRRW EKV YAESHQALVGDKT: AFWLMDKDMY  
 QDEDWRMGDIVHTLTNRRWLEKCVVYAESHQALVGDKTLAFWLMDKDMY  
 ↗590 ↗600 ↗610 ↗620 ↗630  
 ↙650 ↙660 ↙670 ↙680 ↙690  
 DFMALDRPSTSLIDRGIALHKMIRLVTMGLGGEGYLNFMGNEFGHPEWID  
 DFMALDRPST: LIDRGIALHKMIRL: TMGLGGEGYLNFMGNEFGHPEWID  
 DFMALDRPSTPLIDRGIALHKMIRLITMGLGGEGYLNFMGNEFGHPEWID  
 ↗640 ↗650 ↗660 ↗670 ↗680  
 ↙700 ↙710 ↙720 ↙730 ↙740  
 FPRAEOHLSGDSVIPGNQFSYDKCRRRFDLGDAEYLRRLYGLQEFDRPMQY  
 FPR: EOHL: : G: : PGN: SYDKCRRRFDLGDA: YLRY: G: QEFDR: MQ.  
 FPRGEOHLPNGKIVPGNNNSYDKCRRRFDLGDAEYLRRLYGLQEFDRPMQY  
 ↗690 ↗700 ↗710 ↗720 ↗730  
 ↙750 ↙760 ↙770 ↙780 ↙790  
 LEDKYEFTMTSEHQFISRKDEGDRMIVFEKGNLVFVFNFWHTKSYSDYRIA  
 LE: . Y. FMTSEHQ: ISRK: EGDR: I: FE: : NLVFNFWHT: SYSDY: :  
 LEETYGFMTSEHQYISRKNEGDRMIVFERDNLVFNFWHTNSYSDYKVG  
 ↗740 ↗750 ↗760 ↗770 ↗780  
 ↙800 ↙810 ↙820 ↙830 ↙840  
 CLKPGKYKVALDSDDPLFGGFRIDHNAEYFTFEGWYDDRPRSIMVYAPC  
 CLKPGKYK: . LDSDD. LFGGFR: : H. AEYFT FEGWYDDRPRS: : VYAP.  
 CLKPGKYKIVLDSDDTLFGGFNRLNHTAEYFTFEGWYDDRPRSFLVYAPS  
 ↗790 ↗800 ↗810 ↗820 ↗830  
 ↙850 ↙860 ↙870  
 KTAVVYALVDKEEEEEEEEEEEVAA  
 : TAVVYAL. D E. E E. : . V. :  
 RTAVVYALADGVESEPIELSDGVES  
 ↗840 ↗850 ↗860

Fig. 7 SHEET 2

SUBSTITUTE SHEET (RULE 26)

24/75

```

1  -----TTG-AT-----
1  -----TTGA-----
1  -----GA-----
45  AAAAACCTCCTCCACTCAGTCTTGGATCTCTCTCTCTCT

72  TTTCTCTTAATTCCAACCAAGGGAATGAATAAAAGGAT-A
73  TTTCTCTTAATTCCAACCAAGG-AATGAATAAAAGGAT-A
71  TTTCTCTTAATTCCAACCAAGG-AATGAATAAAAGAT-A
165 TTTCTCTTAATTCCAACCAAGG-AATGAATIAAAAGATIA

191 TGTACAAATCTAATGGATTGAGCAGTAATGGTGATCGGAG
191 TGTACAAATCTAATGGATTGAGCAGTAATGGTGATCGGAG
189 TGTACAAATCTAATGGATTGAGCAGTAATGGTGATCGGAG
274 TGTACAAATCTAATGGATTGAGCAGTAATGGTGATCGGAG

311 AATCCGACCTTCTACAGTTGCAGCATCGGGGAAAGTCCT
311 AATCCGACCTTCTACAGTTGCAGCATCGGGGAAAGTCCT
309 AATCCGACCTTCTACAATTGCAGCATCGGGGAAAGTCCT
394 AATCCGACCTTCTACAGTTGCAGCATCGGGGAAAGTCCT

431 CAGCATCAACTGATGTAGATAGTTCAACAATGGAACACGC
431 CAGCATCAACTGATGTAGATAGTTCAACAATGGAACACGC
429 CAGCATCAACTGATGTAGATAGTTCAACAATGGAACACGC
514 CAGCATCAACTGATGTGATAGTTCAACAATGGAACACGC

551 CATCACTACAACCTACAAGAAGGTGGTAACTGGAGGAGTC
551 CATCACTACAACCTACAAGAAGGTGGTAACTGGAGGAGTC
549 CATCACTACAACCTACAAGAAGGTGGTAACTGGAGGAGTC
634 CATCACTACAACCTACAAGAAGGTGGTAACTGGAGGAGTC

671 TTGGTCAGAAGATTTATGAAATAGACCCCCTTTTGACAAA
671 TTGGTCAGAAGATTTATGAAATAGACCCCCTTTTGACAAA
669 TTGGTCAGAAGATTTATGAAATAGACCCCCTTTTGACAAA
754 TTGGTCAGAAGATTTATGAAATAGACCCCCTTTTGACAAA

791 AAGCTTTTCTCGTGGTTATGAAAAAATGGGTTTCACTCG
791 AAGCTTTTCTCGTGGTTATGAAAAAATGGGTTTCACTCG
789 AAGCTTTTCTCGTGGTTATGAAAGAAATGGGTTTCACTCG
874 AAGCTTTTCTCGTGGTTATGAAAAAATGGGTTTCACTCG

```

Fig.8  
Sheet 2

Fig. 8 SHEET 1

SUBSTITUTE SHEET (RULE 26)

25/75

-----GGGCCTTGAAGCTCAGCAATTTGACACTCAGTTAGTTAC  
-----TGGGGCCTTGAAGCTCAGCAATTTGACACTCAGTTAGTTAC  
-----TGGGGCCTTGAAGCTCAGCAATTTGACACTCAGTTAGTTAC  
**TCACGCTTCTCT**TGGGGCCTTGAAGCTCAGCAATTTGACACTCAGTTAGTTAC

GATTTGTAAAAACCTAAGGAGAGAAGAAGAAAGATGGTGTATA**ACTCTCT**  
GATTTGTAAAAACCTAAGGAGAGAAGAAGAAAGATGGTGTATACACTCTCT  
GATTTGTAAAAACCTAAGGAGAGAAGAAGAAAGATGGTGTATACACTCTCT  
GATTTG-----AAGGAGAGAAGAAGAAAGATGGTGTATACACTCTCT

GAATGCTAATGTTTCTGTATTCTTGAAAAAGCACTCTCTTTCACGGAAGATC  
GAATGCTAATGTTTCTGTATTCTTGAAAAAGCACTCTCTTTCACGGAAGATC  
GAATGCTAAT**ATTT**CTGTATTCTTGAAAA**ACTCTCTT**TCACGGAAGATC  
GAATGCTAATGTTTCTGTATTCTTGAAAAAGCACTCTCTTTCACGGAAGATC

TGTGCCTGGA**AG**CCAGAGTGATAGCTCCTCATCCTCAACAGACCAATTTGAG  
TGTGCCTGGA**AG**CCAGAGTGATAGCTCCTCATCCTCAACAGACCAATTTGAG  
TGTGCCTGGAATCCAGAGTGATAGCTCCTCATCCTCAACAG**AT**CAATTTGAG  
TGT**AC**CTGGAATCCAGAGTGATAGCTCCTCATCCTCAACAGACCAATTTGAG

TAGCCAGATTAAAACTGAGAACGATGACGTTGAGCCGTCAAGTGATCTTACA  
TAGCCAGATTAAAACTGAGAACGATGACGTTGAGCCGTCAAGTGATCTTACA  
TAGCCAGATTAAAACTGAGAACGATGACGTTGAGCCGTCAAGTGATCTTACA  
TAGCCAGATTAAAACTGAGAACGATGACGTTGAGCCGTCAAGTGATCTTACA

TAAACATTAAATACTTCTGAAGAGACAATTATTGATGAATCTGATAGGATC  
TAAACATTAAATACTTCTGAAGAGACAATTATTGATGAATCTGATAGGATC  
TAAACATTAAATACTTCTGAAGAGACAATTATTGATGAATCTGATAGGATC  
TAAACATTAAATACTTCTGAAGAGACAATTATTGATGAATCTGATAGGATC

CTATCGTCAACACCTTGATTACAGGTATTCACAGTACAAGAACTGAGGGAG  
CTATCGTCAACACCTTGATTACAGGTATTCACAGTACAAGAACTGAGGGAG  
CTATCGTCAACACCTTGATTACAGGTATTCACAGTACAAGAACTGAGGGAG  
CTATCGTCAACACCTTGATTACAGGTATTCACAGTACAAGAA**AT**GAGGGAG

TAGTGCTACAGGTATCACTTACCGTGAGTGGGCTCCTGGTGCCAGTCAGCT  
TAGTGCTACAGGTATCACTTACCGTGAGTGGGCT**CT**GGTGCCAGTCAGCT  
TAGTGCTACAGGTATCACTTACCGTGAGTGGGCTCCTGGTGCCAGTCAGCT  
TAGTGCTACAGGTATCACTTACCGTGAGTGGGCTCCTGGTGCCAGTCAGCT

Fig. 8  
Sheet  
3

Fig. 8 SHEET 2

SUBSTITUTE SHEET (RULE 26)

26/75

ACTCCTATCACTTATCAGATCTCTATTT 11con.seq  
ACTCCTATCACTTATCAGATCTCTATTT 19con.seq  
ACTCCTATCACTTATCAGATCTCTATTT 10con.seq  
ACTCCTATCACTTATCAGATCTCTATTT psbe2con.seq

GGAGTTCGTTTTCTACTGTTCCATCAG 11con.seq  
GGAGTTCGTTTTCTACTGTTCCATCAG 19con.seq  
GGAGTTCGTTTTCTACTGTTCCATCAG 10con.seq  
GGAGTTCGTTTTCTACTGTTCCATCAG psbe2con.seq

TTGGCTGAAAAGTCTTCTTACAATTCCG 11con.seq  
TTGGCTGAAAAGTCTTCTTACAATTCCG 19con.seq  
TTGGCTGAAAAGTCTTCTTACAATTCCG 10con.seq  
TTGGCTGAAAAGTCTTCTTACAATTCCG psbe2con.seq

TTCACTGAGACATCTCCAGAAAATTCCC 11con.seq  
TTCACTGAGACATCTCCAGAAAATTCCC 19con.seq  
TTCACTGAGACATCTCCAGAAAATTCCC 10con.seq  
TTCACTGAGACATCTCCAGAAAATTCCC psbe2con.seq

GGAAGTGTTGAAGAGCTGGATTTTGCTT 11con.seq  
GGAAGTGTTGAAGAGCTGGATTTTGCTT 19con.seq  
GGAAGTGTTGAAGAGCTGGATTTTGCTT 10con.seq  
GGAAGTGTTGAAGAGCTGGATTTTGCTT psbe2con.seq

AGAGAGAGGGGCATCCCTCCACCTGGAC 11con.seq  
AGAGAGAGGGGCATCCCTCCACCTGGAC 19con.seq  
AGAGAGAGGGGCATCCCTCCACCTGGAC 10con.seq  
AGAGAGAGGGGCATCCCTCCACCTGGAC psbe2con.seq

GCAATTGACAAGTATGAGGGTGGTTTGG 11con.seq  
GCAATTGACAAGTATGAGGGTGGTTTGG 19con.seq  
GCAATTGACAAGTATGAGGGTGGTTTGG 10con.seq  
GCAATTGACAAGTATGAGGGTGGTTTGG psbe2con.seq

CCCCTCATTGGAGATTTCAACAATTGGG 11con.seq  
CCCCTCATTGGAGATTTCAACAATTGGG 19con.seq  
CCCCTCATTGGAGATTTCAACAATTGGG 10con.seq  
CCCCTCATTGGAGATTTCAACAATTGGG psbe2con.seq

Fig. 8  
SHEET 3

SUBSTITUTE SHEET (RULE 26)

27/75

910 ACGCAAATGCTGACATTATGACTCGGAATGAATTTGGTGTC  
911 ACGCAAATGCTGACATTATGACTCGGAATGAATTTGGTGTC  
909 ACGCAAATGCTGACATTATGACTCGGAATGAATTTGGTGTC  
994 ACGCAAATGCTGACATTATGACTCGGAATGAATTTGGTGTC

1030 CTCCATCAGGTGTTAAGGATTCCATTCTGCTTGGATCAAC  
1031 CTCCATCAGGTGTTAAGGATTCCATTCTGCTTGGATCAAC  
1029 CTCCATCAGGTGTTAAGGATTCCATTCTGCTTGGATCAAC  
1114 CTTCATCAGGTGTTAAGGATTCCATTCTGCTTGGATCAAC

1150 AACACCCACGGCCAAAGAAACCAAAGTCGCTGAGAATATAT  
1151 AACACCCACGGCCAAAGAAACCAAAGTCGCTGAGAATATAT  
1149 AACACCCACGGCCAAAGAAACCAAAGTCGCTGAGAATATAT  
1234 AACACCCACGGCCAAAGAAACCAAAGTCGCTGAGAATATAT

1270 TAAAAAA-GCTTGGGTACAATGCGGTGCAATTATGGCTAT  
1271 TAAAAAA-GCTTGGGTACAATGCGGTGCAATTATGGCTAT  
1269 TAAAAAAAGCTTGGGTACAATGCGGTGCAATTATGGCTAT  
1354 TAAAAAACCTTGGGTACAATGCGGTGCAATTATGGCTAT

1389 GACGACCTTAAGTCTTTGATTGATAAAGCTCATGAGCTAGG  
1390 GACGACCTTAAGTCTTTGATTGATAAAGCTCATGAGCTAGG  
1389 GACGACCTTAAGTCTTTGATTGATAAAGCTCATGAGCTAGG  
1473 GACGACCTTAAGTCTTTGATTGATAAAGCTCATGAGCTAGG

1509 GATAGTTGTTACTTTCACTCTGGAGCTCGTGGTTATCATTG  
1510 GATAGTTGTTACTTTCACTCTGGAGCTCGTGGTTATCATTG  
1509 GATAGTTGTTACTTTCACTCTGGAGCTCGTGGTTATCATTG  
1593 GATAGTTGTTACTTTCACTCTGGAGCTCGTGGTTATCATTG

1628 GATGAGTTCAAATTTGATGGATTTAGATTTGATGGTGTGAC  
1630 GATGAGTTCAAATTTGATGGATTTAGATTTGATGGTGTGAC  
1629 GATGAGTTCAAATTTGATGGATTTAGATTTGATGGTGTGAC  
1713 GATGAGTTCAAATTTGATGGATTTAGATTTGATGGTGTGAC

1748 GTGGATGCTGTTGTGTATCTGATGCTGGTCAACGATCTTAT  
1750 GTGGATGCTGTTGTGTATCTGATGCTGGTCAACGATCTTAT  
1749 GTGGATGCTGTTGTGTATCTGATGCTGGTCAACGATCTTAT  
1833 GTGGATGCTGTTGTGTATCTGATGCTGGTCAACGATCTTAT

Fig. 8  
Sheet 5Fig. 8  
SHEET 4

SUBSTITUTE SHEET (RULE 26)

28/75

TGGGAGATTTTTCTGCCAAATAATGTGGATGGTTCTCCTGCAATTC  
TGGGAGATTTTTCTGCCAAATAATGTGGATGGTTCTCCTGCAATTC  
TGGGAGATTTTTCTGCCAAATAATGTGGATGGTTCTCCTGCAATTC  
TGGGAGATTTTTCTGCCAAATAATGTGGATGGTTCTCCTGCAATTC

TACTCTTTACAGCTTCCTGATGAAATTCCATATAATGGAATATATT  
TACTCTTTACAGCTTCCTGATGAAATTCCATATAATGGAATATT  
TACTCTTTACAGCTTCCTGATGAAATTCCATATAATGGAATATATT  
TACTCTTTACAGCTTCCTGATGAAATTCCATATAATGGAATATATT

GAATCTCATATTGGAATGAGTAGTCCGGAGCCTAAAATTAACATCAT  
GAATCTCATATTGGAATGAGTAGTCCGGAGCCTAAAATTAACATCAT  
GAATCTCATATTGGAATGAGTAGTCCGGAGCCTAAAATTAACATCAT  
GAATCTCATATTGGAATGAGTAGTCCGGAGCCTAAAATTAACATCAT

TCAAGAGCATTCTTATTATGCTAGTTTTGGTTATCATGTCACAAAT  
TCAAGAGCATTCTTATTATGCTAGTTTTGGTTATCATGTCACAAAT  
TCAAGAGCATTCTTATTATGCTAGTTTTGGTTATCATGTCACAAAT  
TCAAGAGCATTCTTATTATGCTAGTTTTGGTTATCATGTCACAAAT

AATTGTTGTTCTCATGGACATTGTTACAGCCATGCATCAAATAAT  
AATTGTTGTTCTCATGGACATTGTTACAGCCATGCATCAAATAAT  
AATTGTTGTTCTCATGGACATTGTTACAGCCATGCATCAAATAAT  
AATTGTTGTTCTCATGGACATTGTTACAGCCATGCATCAAATAAT

GATGTGGGATTCCGCCTCTTTAACTATGGAACTGGGAGGTACTT  
GATGTGGGATTCCGCCTCTTTAACTATGGAACTGGGAGGTACTT  
GATGTGGGATTCCGCCTCTTTAACTATGGAACTGGGAGGTACTT  
GATGTGGGATTCCGCCTCTTTAACTATGGAACTGGGAGGTACTT

ATCAATGATGTATACTCACCACGGATTATCGGTGGGATTCACTGGG  
ATCAATGATGTATACTCACCACGGATTATCGGTGGGATTCACTGGG  
ATCAATGATGTATACTCACCACGGATTATCGGTGGGATTCACTGGG  
ATCAATGATGTATACTCACCACGGATTATCGGTGGGATTCACTGGG

TCATGGGCTTTTCCCAGATGCAATTACCATTGGTGAAGATGTTAGC  
TCATGGGCTTTTCCCAGATGCAATTACCATTGGTGAAGATGTTAGC  
TCATGGGCTTTTCCCAGATGCAATTACCATTGGTGAAGATGTTAGC  
TCATGGGCTTTTCCCAGATGCAATTACCATTGGTGAAGATGTTAGC

Fig. 8  
Sheet 6

Fig. 8  
SHEET 5

SUBSTITUTE SHEET (RULE 26)

29/75

CTCATGGGTCCAGAGTGAAGATACGTATGGACA 11con.seq  
 CTCATGGGTCCAGAGTGAAGATACGTATGGACA 19con.seq  
 CTCATGGGTCCAGAGTGAAGATACGTATGGACA 10con.seq  
 CTCATGGGTCCAGAGTGAAGATACGATGGACA psbe2con.seq

ATGATCCACCCGAAGAGGAGAGGTATATCTTCC 11con.seq  
 ATGATCCACCCGAAGAGGAGAGGTATATCTTCC 19con.seq  
 ATGATCCACCCGAAGAGGAGAGGTATATCTTCC 10con.seq  
 ATGATCCACCCGAAGAGGAGAGGTATCTCTTCC psbe2con.seq

ACGTGAATTTTAGAGATGAAGTTCTTCCTCGCA 11con.seq  
 ACGTGAATTTTAGAGATGAAGTTCTTCCTCGCA 19con.seq  
 ACGTGAATTTTAGAGATGAAGTTCTTCCTCGCA 10con.seq  
 ACGTGAATTTTAGAGATGAAGTTCTTCCTCGCA psbe2con.seq

TTTTTGCACCAAGCAGCCGTTTTGGAACGCC 11con.seq  
 TTTTTGCACCAAGCAGCCGTTTTGGAACGCC 19con.seq  
 TTTTTGCACCAAGCAGCCGTTTTGGAACGCC 10con.seq  
 TTTTTGCACCAAGCAGCCGTTTTGGAACGCC psbe2con.seq

ACTTTAGATGGACTGAACATGTTTGACGGCACC 11con.seq  
 ACTTTAGATGGACTGAACATGTTTGACGGCACC 19con.seq  
 ACTTTAGATGGACTGAACATGTTTGACGGCACA 10con.seq  
 ACTTTAGATGGACTGAACATGTTTGACGGCACA psbe2con.seq

AGGTATCTTCTCTCAAATGCGAGATGGTGGTTG 11con.seq  
 AGGTATCTTCTCTCAAATGCGAGATGGTGGTTG 19con.seq  
 AGGTATCTTCTCTCAAATGCGAGATGGTGGTTG 10con.seq  
 AGGTATCTTCTCTCAAATGCGAGATGGTGGTTG psbe2con.seq

AACTACGAGGAATACTTTGGACTCGCAACTGAT 11con.seq  
 AACTACGAGGAATACTTTGGACTCGCAACTGAT 19con.seq  
 AACTACGAGGAATACTTTGGACTCGCAACTGAT 10con.seq  
 AACTACGAGGAATACTTTGGACTCGCAACTGAT psbe2con.seq

GGAATGCCGACATTTTGTATTCCCGTTCAAGAT 11con.seq  
 GGAATGCCGACATTTTGTATTCCCGTTCAAGAT 19con.seq  
 GGAATGCCGACATTTTGTATTCCCGTTCAAGAT 10con.seq  
 GGAATGCCGACATTTTGTATTCCCGTTCAAGAT psbe2con.seq

Fig. 8  
 SHEET 6

SUBSTITUTE SHEET (RULE 26)





31/75

TGATAAATGGATTGAGTTGCTCAAGAAACGGGATGAGGATTGGAGA  
TGATAAA[GGATTGAGTTGCTCAAGAAACGGGATGAGGATTGGAGA  
TGATAAATGGATTGAGTTGCTCAAGAAACGGGATGAGGATTGGAGA  
TGATAAATGGATTGAGTTGCTCAAGAAACGGGATGAGGATTGGAGA

TCAAGCTCTAGTCGGTGATAAACTATAGCATTCTGGCTGATGGAC  
TCAAGCTCTAGTCGGTGATAAACTATAGCATTCTGGCTGATGGAC  
TCAAGCTCTAGTCGGTGATAAACTATAGCATTCTGGCTGATGGAC  
TCAAGCTCTAGTCGGTGATAAACTATAGCATTCTGGCTGATGGAC

GATGATTAGGCTTGTAACCTATGGGATTAGGAGGAGAAGGGTACCTA  
GATGATTAGGCTTGTAACCTATGGGATTAGGAGGAGAAGGGTACCTA  
GATGATTAGGCTTGTAACCTATGGGATTAGGAGGAGAAGGGTACCTA  
GATGATTAGGCTTGTAACCTATGGGATTAGGAGGAGAAGGGTACCTA

CTCAGTAATTCCTCGGAAACCAATTCAGTTATGATAAATGCAGACGG  
CTCAGTAAT[CTCGGAAACCAATTCAGTTATGATAAATGCAGACGG  
CTCAGTAATTCCT[AGAAACCAATTCAGTTATGATAAATGCAGACGG  
CTCAGTAATTCCTCGGAAACCAATTCAGTTATGATAAATGCAGACGG

TGAAGATAAATATGAGTTTATGACTTCAGAACACCAGTTCATATCA  
TGAAGATAAATATGAGTTTATGACTTCAGAACACCAGTTCATATCA  
TGAAGATAAATATGAGTTTATGACTTCAGAACACCAGTTCATATCA  
TGAAGATAAATATGAGTTTATGACTTCAGAACACCAGTTCATATCA

AAA[AGCTATTCAGACTATCGCATAGGCTGCCTGAAGCCTGGAAAA  
AAAAAGCTATTCAGACTATCGCATAG[CTGCCTGAAGCCTGGAAAA  
AAAA[AGCTATTCAGACTATCGCATAGGCTGCCTGAAGCCTGGAAAA  
AAAAAGCTATTCAGACTATCGCATAGGCTG[CTGAAGCCTGGAAAA

CACCT[CTGAAGGAT[GTATGATGATCGTCCT[GTTCATTATGGTG  
CACCTTTGAAGGATGGTATGATGATCGTCCTCGTTCAATTATGGTG  
CACCTTTGAAGGATGGTATGATGATCGTCCTCGTTCAATTATGGTG  
CACCTTTGAAGGATGGTATGATGATCGTCCTCGTTCAATTATGGTG

-----TAGCAGTAGTAGAAGAA[CCCATTG-----AAGAATGAACG  
AGAAGTAGCAG[AGTAGAAGAAGTAGTAGTAGAAGAAGAATGAACG  
-----TAGCAGTAGTAGAAGAAGTAGTAGTAGAAGAAGAATGAACG  
-----TAGCAGTAGTAGAAGAAGTAGTAGTAGAAGAAGAATGAACG

Fig. 8  
Sheet 9

Fig. 8  
SHEET 8

SUBSTITUTE SHEET (RULE 26)

32/75

GTGGGTGATATTGTTTCATACACTGACAAATAGA 11con.seq  
GTGGGTGATATTGTTTCATACACTGACAAATAGA 19con.seq  
GTGGGTGATATTGTTTCATACACTGACAAATAGA 10con.seq  
GTGGGTGATATTGTTTCATACACTGACAAATAGA psbe2con.seq

AAGGATATGTATGATTTTATGGCTCTGGATAGA 11con.seq  
AAGGATATGTATGATTTTATGGCTCTGGATAGA 19con.seq  
AAGGATATGTATGATTTTATGGCTCTGGATAGA 10con.seq  
AAGGATATGTATGATTTTATGGCTTTGGATAGA psbe2con.seq

AATTTTCATGGGAAATGAATTCGGCCACCCTGAG 11con.seq  
AATTTTCATGGGAAATGAATTCGGCCACCCTGAG 19con.seq  
AATTTTCATGGGAAATGAATTCGGCCACCCTGAG 10con.seq  
AATTTTCATGGGAAATGAATTCGGCCACCCTGAG psbe2con.seq

AGATTTGACCTGGGAGATGCAGAATATTTAAGA 11con.seq  
AGATTTGACCTGGGAGATGCAGAATATTTAAGA 19con.seq  
AGATTTGACCTGGGAGATGCAGAATATTTAAGA 10con.seq  
AGATTTGACCTGGGAGATGCAGAATATTTAAGA psbe2con.seq

CGAAAGGATGAAGGAGATAGGATGATTGTATTT 11con.seq  
CGAAAGGATGAAGGAGATAGGATGATTGTATTT 19con.seq  
CGAAAGGATGAAGGAGATAGGATGATTGTATTT 10con.seq  
CGAAAGGATGAAGGAGATAGGATGATTGTATTT psbe2con.seq

TACAAGGTTGCTTGGACTCAGATGATCCACTT 11con.seq  
TACAAGGTTGCCTTGGACTCAGATGATCCACTT 19con.seq  
TACAAGGTTGCCTTGGACTCAGATGATCCACTT 10con.seq  
TACAAGGTTGCCTTGGACTCAGATGATCCACTT psbe2con.seq

TATGCACCTAGTAGAACAGCAGTGGTCTATGCA 11con.seq  
TATGCACCTTGTAAACAGCAGTGGTCTATGCA 19con.seq  
TATGCACCTAGTAGAACAGCAGTGGTCTATGCA 10con.seq  
TATGCACCTAGTAGAACAGCAGTGGTCTATGCA psbe2con.seq

AACTTGTGATCGCGTTGAAAGATTTGAACGTTA 11con.seq  
AACTTGTGATCGCGTTGAAAGATTTGAACG--- 19con.seq  
AACTTGTGATCGCGTTGAAAGATTTGAACG--- 10con.seq  
AACTTGTGATCGCGTTGAAAGATTTGAACG--- psbe2con.seq

Fig. 8  
SHEET 9

SUBSTITUTE SHEET (RULE 26)

33/75

2795 CTTGGTCATCCACATAGAGCTTCTTGAC-----  
2827 -----CTACATAGAGCTTCTTGACGTATCTGGCAATAT  
2814 -----CCACATAGAGCTTCTTGACGTATCTGGCAATAT  
2895 -----CTACATAGAGCTTCTTGACGTATCTGGCAATAT  
  
2898 AGAGATGAAGTGCTGAACAAA--CATATGTAAAATCGATGAA  
2937 AGAGATGAAGTGCTGAACAAA--CATATGTAAAATCGATGAA  
2924 AGAGATGAAGTGCTGAACAAAACATATGTAAAATCGATGAA  
3005 AGAGATGAAGTGCTGAACAAA--CATATGTAAAATCGATGAA  
  
2975  
3012  
3003  
3123 GCCCACTAGAAATCAATTATGTGAGACCTAAAAACAATAAC

Fig. 8  
Sheet 11

Fig. 8 SHEET 10

SUBSTITUTE SHEET (RULE 26)

34/75

---ATCAGTCTTGGCGGAATTTCATGTGACAAAGGTTTGCACTT  
TGCATCAGTCTTGGCGGAATTTTCATGTGACAC-AAGGTTTGCAATT  
TGCATTAGTCTTGGCGGAATTTTCATGTGACAA-CAGGTTTGCAATT  
TGCATCAGTCTTGGCGGAATTTTCATGTGACAA-AAGGTTTGCAATT

TTTATGTCGAATGCTGGGACGATCGAATTCCTGCAGCC  
TTTATGTCGAATGCTGGGACGATCGAATTCCTGCAG  
TTTATGTCGAATGCTGGGACGATCGAATTCCTGCAGCC  
TTTATGTCGAATGCTGGGACGGCTTCAGCAGTTTGCTTAGTGA

Fig. 8  
Sheet 12

CATAAAATGGAAATAGTGCTGATCTAATGATGTTTAAANCCNNNA

Fig. 8 SHEET 11

SUBSTITUTE SHEET (RULE 26)

35/75

---

CTTTCCACTATTAGTAGTCAACGATATACGC 11con.seq  
CTTTCCACTATTAGTAGTGCAACGATATACGC 19con.seq  
CTTTCCACTATTAGTAGTGCAACGATATACGC 10con.seq  
CTTTCCACTATTAGTAGTGCAACGATATACGC psbe2con.seq

11con.seq

19con.seq

10con.seq

GTTCTGTAAATTGTCATCTCTTTANATGTACA psbe2con.seq

11con.seq

19con.seq

10con.seq

AAAAAAAAAAAAAACTCGAG

psbe2con.seq

Fig. 8 SHEET 12

36/75

GGATGCTAATGTTTCTGTATTCTTGAAAAAGCACTCTCTTTCACGG  
CCTACGATTACAAAGACATAAGAACTTTTTCTGTGAGAGAAAGTGCC  
A N V S V F L K K H S L S R

TTCTACAGTTGCAGCATCGGGGAAAGTCCTTGTGCCTGGAAYCCAG  
AAGATGTCAACGTCGTAGCCCCCTTTCAGGAACACGGACCTTRGGTC  
S T V A A S G K V L V P G ? Q

GACATCTCCAGAAAATTCCCCAGCATCAACTGATGTAGATAGTTCA  
CTGTAGAGGTCTTTTAAGGGGTCGTAGTTGACTACATCTATCAAGT  
T S P E N S P A S T D V D S S

TGAGCCGTCAAGTGATCTTACAGGAAGTGTTGAAGAGCTGGATTTT  
ACTCGGCAGTTCCTAGAAATGTCCTTCACAACTTCTCGACCTAAAA  
E P S S D L T G S V E E L D F

TAAACATTAAATACTTCTGAAGAGACAATTATTGATGAATCTGAT  
ATTTTGTAATTTATGAAGACTTCTCTGTTAATAACTACTTAGACTA  
K T L N T S E E T I I D E S D

Hinc II  
GATTTATGAAATAGACCCCCTTTTGACAACTATCGTCAACACCTT  
CTAAATACTTTATCTGGGGGAAAAGTGTGATAGCAGTTGTGGAA  
I Y E I D P L L T N Y R Q H L

Fig.9  
Sheet  
2

Fig.9 SHEET 1

SUBSTITUTE SHEET (RULE 26)

37/75

Bgl II

AAGATCTTGGCTGAAAAGTCTTCTTACAATTCCGAATCCCGACC

TTCTAGAACCGACTTTTTCAGAAGAATGTTAAGGCTTAGGGCTGG

K I L A E K S S Y N S E S R P

90

AGTGATAGCTCCTCATCCTCAACAGACCAATTTGAGTTCACTGA

TCACTATCGAGGAGTAGGAGTTGTCTGGTTAACTCAAGTGACT

S D S S S S S T D Q F E F T E

180

ACAATGGAACACGCTAGCCAGATTAAAACTGAGAACGATGACGT

TGTTACCTTGTGCGATCGGTCTAATTTTGACTCTTGCTACTGCA

T M E H A S Q I K T E N D D V

270

GCTTCATCACTACAACCTACAAGAAGGTGGTAACTGGAGGAGTC

CGAAGTAGTGATGTTGATGTTCTTCCACCATTTGACCTCCTCAG

A S S L Q L Q E G G K L E E S

360

AGGATCAGAGAGAGGGGCATCCCTCCACCTGGACTTGGTCAGAA

TCCTAGTCTCTCTCCCCGTAGGGAGGTGGACCTGAACCAGTCTT

R I R E R G I P P P G L G Q K

450

GATTACAGGTATTCACAGTACAAGAACTGAGGGAGGCAATTGA

CTAATGTCCATAAGTGTCATGTTCTTTGACTCCCTCCGTAACT

D Y R Y S Q Y K K L R E A I D

540

Fig. 9 SHEET 2

SUBSTITUTE SHEET (RULE 26)

38/75

Hind III

CAAGTATGAGGGTGGTTTGGAGCTTTTTCTCGTGGTTATGAAAAA  
GTTCATACTCCCAACCAACCTTCGAAAAAGAGCACCAATACTTTTT  
K Y E G G L E A F S R G Y E K

Pvu II

GGCTCCTGGTGCCAGTCAGCTGCCCTCATTGGAGATTTCAACAAT  
CCGAGGACCACGGGTCAGTCGACGGGAGTAACCTCTAAAGTTGTTA  
A P G A Q S A A L I G D F N N

CTGGGAGATTTTTCTGCCAAATAATGTGGATGGTTCTCCTGCAATT  
GACCCTCTAAAAAGACGGTTTATTACACCTACCAAGAGGACGTAA  
W E I F L P N N V D G S P A I

TGTTAAGGATTCCATTCTGCTTGGATCAACTACTCTTTACAGCTT  
ACAATTCCTAAGGTAAGGACGAACCTAGTTGATGAGAAATGTCGAA  
V K D S I P A W I N Y S L Q L

AGAGGAGAGGTATRTCTTCCAACACCCACGGCCAAAGAAACCAAAG  
TCTCCTCTCCATAYAGAAGTTGTGGGTGCCGGTTTCTTTGGTTTC  
E E R Y ? F Q H P R P K K P K

Fig.9  
Sheet  
4

Fig.9 SHEET 3

SUBSTITUTE SHEET (RULE 26)



39/75

ATGGGTTTCACTCGTAGTGCTACAGGTATCACTTACCGTGAGTG  
TACCCAAAGTGAGCATCACGATGTCCATAGTGAATGGCACTCAC  
M G F T R S A T G I T Y R E W 630

TGGGACGCAAATGCTGACATTATGACTCGGAATGAATTTGGTGT  
ACCCTGCGTTTACGACTGTAATACTGAGCCTTACTTAAACCACA  
W D A N A D I M T R N E F G V 720

CCTCATGGGTCCAGAGTGAAGATACGYATGGACACTCCATCAGG  
GGAGTACCCAGGTCTCACTTCTATGCRTACCTGTGAGGTAGTCC  
P H G S R V K I R M D T P S G 810

CCTGATGAAATTCCATATAATGGAATATATTATGATCCACCCGA  
GGACTACTTTAAGGTATATTACCTTATATAATACTAGGTGGGCT  
P D E I P Y N G I Y Y D P P E 900

TCGCTGAGAATATATGAATCTCATATTGGAATGAGTAGTCCGGA  
AGCGACTCTTATATACTTAGAGTATAACCTTACTCATCAGGCCT  
S L R I Y E S H I G M S S P E 990

Fig. 9 SHEET 4

SUBSTITUTE SHEET (RULE 26)

40/75

Xmn I

GCCTAAAATTAACATCATACGTGAATTTTAGAGATGAAGTTCTTCCT  
CGGATTTTAATTGAGTATGCACTTAAAATCTCTACTTCAAGAAGGA  
P K I N S Y V N F R D E V L P

TCAAGAGCATTCTTATTATGCTAGTTTTGGTTATCATGTCACAAAT  
AGTTCTCGTAAGAATAATACGATCAAACCAATAGTACAGTGTTTA  
Q E H S Y Y A S F G Y H V T N

GTCTTTGATTGATAAAGCTCATGAGCTAGGAATTGTTGTTCTCATG  
CAGAACTAACTATTTTCGAGTACTCGATCCTTAACAACAAGAGTAC  
S L I D K A H E L G I V V L M

GAACATGTTTGACGGCACAGATAGTTGTTACTTTCACTCTGGAGCT  
CTTGTACAACTGCCGTGTCTATCAACAATGAAAGTGAGACCTCGA  
N M F D G T D S C Y F H S G A

AACTGGGAGGTACTTAGGTATCTTCTCTCAAATGCGAGATGGTGG  
TTTGACCCTCCATGAATCCATAGAAGAGAGTTTACGCTCTACCACC  
N W E V L R Y L L S N A R W W

ATCAATGATGTATACTCACCACGGATTATCGGTGGGATTCACTGGG  
TAGTTACTACATATGAGTGGTGCCTAATAGCCACCCTAAGTGACCC  
S M M Y T H H G L S V G F T G

Fig.9  
Sheet  
6

Fig.9 SHEET 5

SUBSTITUTE SHEET (RULE 26)

41/75

CGCATAAAAAASCTTGGGTACAATGCGGTGCAAATTATGGCTAT  
+-----+ 1080  
GCGTATTTTTTSGAACCCATGTTACGCCACGTTTAATACCGATA  
R I K ? L G Y N A V Q I M A I

TTTTTTGCACCAAGCAGCCGTTTTTGAACGCCCCGACGACCTTAA  
+-----+ 1170  
AAAAAACGTGGTTCGTCGGCAAACCTTGCGGGCTGCTGGAATT  
F F A P S S R F G T P D D L K

GACATTGTTACAGCCATGCATCAAATAATACTTTAGATGGACT  
+-----+ 1260  
CTGTAACAAGTGTCGGTACGTAGTTTATTATGAAATCTACCTGA  
D I V H S H A S N N T L D G L

Sac I

CGTGGTTATCATTGGATGTGGGATTCCCGCCTCTTTAACTATGG  
+-----+ 1350  
GCACCAATAGTAACCTACACCCTAAGGGCGGAGAAATTGATACC  
R G Y H W M W D S R L F N Y G

TTGGATGAGTTCAAATTTGATGGATTTAGATTTGATGGTGTGAC  
+-----+ 1440  
AACCTACTCAAGTTTAAACTACCTAAATCTAAACTACCACTG  
L D E F K F D G F R F D G V T

AACTACGAGGAATACTTTGGACTCGCAACTGATGTGGATGCTGT  
+-----+ 1530  
TTGATGCTCCTTATGAAACCTGAGCGTTGACTACACCTACGACA  
N Y E E Y F G L A T D V D A V

Fig. 9 SHEET 6

SUBSTITUTE SHEET (RULE 26)

42/75

Hinc II

TGTGTATCTGATGCTGGTCAACGATCTTATTCACGGGCTTTTCCCA  
ACACATAGACTACGACCAGTTGCTAGAATAAGTGCCCGAAAAGGGT  
V Y L M L V N D L I H G L F P

TTGTATTCCCGTTCAAGATGGGGGTGTTGGCTTTGACTATCGGCTG  
AACATAAGGGCAAGTTCTACCCCCACAACCGAAACTGATAGCCGAC  
C I P V Q D G G V G F D Y R L

GGATGAGGATTGGAGAGTGGGTGATATTGTTTCATACACTGACAAAT  
CCTACTCCTAACCTCTCACCCACTATAACAAGTATGTGACTGTTTA  
D E D W R V G D I V H T L T N

TCAAGCTCTAGTCGGTGATAAAACTATAGCATYCTGGCTGATGGAC  
AGTTTCGAGATCAGCCACTATTTTGATATCGTARGACCGACTACCTG  
Q A L V G D K T I A ? W L M D

ATTAATAGATCGTGGGATAGCATTGCACAAGATGATTAGGCTTGTA  
TAATTATCTAGCACCTATCGTAACGTGTTCTACTAATCCGAACAT  
L I D R G I A L H K M I R L V

Fig.9  
Sheet  
8

Fig. 9 SHEET 7

SUBSTITUTE SHEET (RULE 26)

43/75

GATGCAATTACCATTGGTGAAGATGTTAGCGGAATGCCGACATT 1620  
CTACGTTAATGGTAACCACTTCTACAATCGCCTTACGGCTGTAA  
D A I T I G E D V S G M P T F

Nde I

CATATGGCAATTGCTGATAAATGGATTGAGTTGCTCAAGAAACG 1710  
GTATACCGTTAACGACTATTTACCTAACTCAACGAGTTCTTTGC  
H M A I A D K W I E L L K K R

AGAAGATGGTCGGAAAAGTGTGTTTCATMCGCTGAAAGTCATGA 1800  
TCTTCTACCAGCCTTTTCACACAAAGTAKGCGACTTTTCAGTACT  
R R W S E K C V S ? A E S H D

Hinc II

AAGGATATGTATGATTTTATGGCTCTGGATAGACCGTCAACATC 1890  
TTCCTATACATACTAAAATACCGAGACCTATCTGGCAGTTGTAG  
K D M Y D F M A L D R P S T S

Asp 718

Kpn I

ACTATGGGATTAGGAGGAGAAGGGTACCTAAATTTTCATGGGAAA 1980  
TGATACCCTAATCCTCCTCTTCCCATGGATTAAAGTACCCTTT  
T M G L G G E G Y L N F M G N

Fig. 9 SHEET 8

SUBSTITUTE SHEET (RULE 26)

44/75

EcoR I

TGAATTCGGCCACCCTGAGTGGATTGATTTCCTAGGGCTGARCAA  
ACTTAAGCCGGTGGGACTCACCTAACTAAAGGGATCCCGACTYGT  
E F G H P E W I D F P R A E Q

Ssp I

TGATAAATGCAGACGGAGATTTGACCTGGGAGATGCAGAATATTTA  
ACTATTTACGTCTGCCTCTAACTGGACCCTCTACGTCTTATAAAT  
D K C R R R F D L G D A E Y L

TGAAGATAAATATGAGTTTATGACTTCAGAACACCAGTTCATATCA  
ACTTCTATTTATACTCAAATACTGAAGTCTTGTGGTCAAGTATAGT  
E D K Y E F M T S E H Q F I S

CCTAGTTTTTGTCTTTAATTTTCACTGGACAAATAGCTATTCAGAC  
GGATCAAAAACAGAAATTAAGTGACCTGTTTATCGATAAGTCTG  
L V F V F N F H W T N S Y S D

GGACTCAGATGATCCACTTTTTGGTGGCTTCGGGAGAATTGATCAT  
CCTGAGTCTACTAGGTGAAAAACCACCGAAGCCCTCTTAAGTAGTA  
D S D D P L F G G F G R I D H

YCGYYCAATTATGGTGTATGCACCTAGTAGAACAGCAGTGGTCTAT  
RGCRRGTTAATACCACATACGTGGATCATCTTGTGTCACCAGATA  
R ? I M V Y A P S R T A V V Y

NGAAGAATTTT  
NCTTCTTAAAA  
E E F

2531

Fig 9 SHEET 9

SUBSTITUTE SHEET (RULE 26)

Fig 9  
Sheet  
10

45/75

CACCTCTCTGATGGCTCAGTAATTCCCGGAAACCAATTCAGTTA  
GTGGAGAGACTACCGAGTCATTAAGGGCCTTTGGTTAAGTCAAT 2070  
H L S D G S V I P G N Q F S Y

Nco I

AGATACCATGGGTTGCAAGAATTTGACCGGGCTATGCAGTATCT  
TCTATGGTACCCAACGTTCTTAAACTGGCCCGATACGTCATAGA 2160  
R Y H G L Q E F D R A M Q Y L

CGAAAGGATGAAGGAGATAGGATGATTGTATTTGAAARAGGAAA  
GCTTTCCTACTTCCTCTATCCTACTAACATAAACTTTTTCCTTT 2250  
R K D E G D R M I V F E ? G N

TATCGCATAGGCTGCCTGAAGCCTGGAAAATACAAGGTTGGCTT  
ATAGCGTATCCGACGGACTTCGGACCTTTTATGTTCCAACCGAA 2340  
Y R I G C L K P G K Y K V G L

Ssp I

AATGCCGAATATTTACCTCTGAAGGATCGTATGATGATCGYCC  
TTACGGCTTATAAAGTGGAGACTTCCTAGCATACTACTAGCRGG 2430  
N A E Y F T S E G S Y D D R P

GCACTAGTAGACAAANTAGAAGNAGAAGAAGAAGAANCCGN  
CGTGATCATCTGTTTATCTTCNTCTTCTTCTTCTTNGGCN 2520  
A L V D K ? E ? E E E E ? ?

Fig. 9 SHEET 10

SUBSTITUTE SHEET (RULE 26)


Fig.10  
Sheet 2

Fig. 10 SHEET 1

SUBSTITUTE SHEET (RULE 26)



47/75

40	50	60	70	
TAGTTACACT	CC	ATCACTTATCAGATCTCTAT		10con. seq
TAGTTACACT	CCTATCACTTATCAGATCTCTAT			11con. seq
TAGTTACACT	CCTATCACTTATCAGATCTCTAT			19con. seq
	-----CATT	-----		86CON. SEQ
	-----	-----		pcrsbe2con. seq
110	120	130	140	
GATAGATTTGTAAAAACCCTAAGGAGAGAAGAA				10con. seq
GATAGATTTGTAAAAACCCTAAGGAGAGAAGAA				11con. seq
GATAGATTTGTAAAAACCCTAAGGAGAGAAGAA				19con. seq
GAGAAATT	-----AACTATCAGAGGA	-----		86CON. SEQ
	-----	-----		pcrsbe2con. seq
180	190	200	210	
TACTGTTCCATCAGTGTACAAATCTAATGGATT				10con. seq
TACTGTTCCATCAGTGTACAAATCTAATGGATT				11con. seq
TACTGTTCCATCAGTGTACAAATCTAATGGATT				19con. seq
CACCAT	-----CACCA	-----T		86CON. SEQ
	-----	-----		pcrsbe2con. seq
250	260	270	280	
GTATTCTTGAAAAA	ACACTCTCTTTACGGAAG			10con. seq
GTATTCTTGAAAAAGCACTCTCTTTACGGAAG				11con. seq
GTATTCTTGAAAAAGCACTCTCTTTACGGAAG				19con. seq
	-----CCATGG	-----G		86CON. SEQ
	-----	-----		pcrsbe2con. seq
320	330	340	350	
GACCTTCTACAA	ATTGCAGCATCGGGGAAAGTCC			10con. seq
GACCTTCTACAGTTGCAGCATCGGGGAAAGTCC				11con. seq
GACCTTCTACAGTTGCAGCATCGGGGAAAGTCC				19con. seq
GACCTTCTACAGTTGCAGCATCGGGGAAAGTCC				86CON. SEQ
GACCTTCTACAGTTGCAGCATCGGGGAAAGTCC				pcrsbe2con. seq

Fig. 10 SHEET 2

SUBSTITUTE SHEET (RULE 26)

48/75

	360	370	380
348	TTGTGCCTGGAAT	CCAGAGTGATAGCTCCTCATCCTC	
350	TTGTGCCTGGAACCCAGAGTGATAGCTCCTCATCCTC		
350	TTGTGCCTGGAACCCAGAGTGATAGCTCCTCATCCTC		
127	TTGTGCCTGGAACCCAGAGTGATAGCTCCTCATCCTC		
120	TTGTGCCTGGAAT	CCAGAGTGATAGCTCCTCATCCTC	
	430	440	450
418	AGAAAATTCCCCAGCATCAACTGATGTAGATAGTTCA		
420	AGAAAATTCCCCAGCATCAACTGATGTAGATAGTTCA		
420	AGAAAATTCCCCAGCATCAACTGATGTAGATAGTTCA		
197	AGAAAATTCCCCAGCATCAACTGATGTAGATAGTTCA		
190	AGAAAATTCCCCAGCATCAACTGATGTAGATAGTTCA		
	500	510	520
488	AACGATGACGTTGAGCCGTCAAGTGATCTTACAGGAA		
490	AACGATGACGTTGAGCCGTCAAGTGATCTTACAGGAA		
490	AACGATGACGTTGAGCCGTCAAGTGATCTTACAGGAA		
267	AACGATGACGTTGAGCCGTCAAGTGATCTTACAGGAA		
260	AACGATGACGTTGAGCCGTCAAGTGATCTTACAGGAA		
	570	580	590
558	AACTACAAGAAGGTGGTAAACTGGAGGAGTCTAAAAC		
560	AACTACAAGAAGGTGGTAAACTGGAGGAGTCTAAAAC		
560	AACTACAAGAAGGTGGTAAACTGGAGGAGTCTAAAAC		
337	AACTACAAGAAGGTGGTAAACTGGAGGAGTCTAAAAC		
330	AACTACAAGAAGGTGGTAAACTGGAGGAGTCTAAAAC		
	640	650	660
628	ATCTGATAGGATCAGAGAGAGGGGCATCCCTCCACCT		
630	ATCTGATAGGATCAGAGAGAGGGGCATCCCTCCACCT		
630	ATCTGATAGGATCAGAGAGAGGGGCATCCCTCCACCT		
407	ATCTGATAGGATCAGAGAGAGGGGCATCCCTCCACCT		
400	ATCTGATAGGATCAGAGAGAGGGGCATCCCTCCACCT		

Fig.10  
Sheet 4

Fig.10 SHEET 3

SUBSTITUTE SHEET (RULE 26)

49/75

390	400	410	420	
AACAGATCAATTTGAGTTCCTGAGACATCTCC				10con. seq
AACAGACCAATTTGAGTTCACTGAGACATCTCC				11con. seq
AACAGACCAATTTGAGTTCACTGAGACATCTCC				19con. seq
AACAAACCAATTTGAGTTCACTGAGACATCTCC				86CON. SEQ
AACAGACCAATTTGAGTTCACTGAGACATCTCC				pcrsbe2con. seq
460	470	480	490	
ACAATGGAACACGCTAGCCAGATTAAAACTGAG				10con. seq
ACAATGGAACACGCTAGCCAGATTAAAACTGAG				11con. seq
ACAATGGAACACGCTAGCCAGATTAAAACTGAG				19con. seq
ACAATGGAACACGCTAGCCAGATTAAAACTGAG				86CON. SEQ
ACAATGGAACACGCTAGCCAGATTAAAACTGAG				pcrsbe2con. seq
530	540	550	560	
GTGTTGAAGAGCTGGATTTTGCTTCATCACTAC				10con. seq
GTGTTGAAGAGCTGGATTTTGCTTCATCACTAC				11con. seq
GTGTTGAAGAGCTGGATTTTGCTTCATCACTAC				19con. seq
GTGTTGAAGAGCTGGATTTTGCTTCATCACTAC				86CON. SEQ
GTGTTGAAGAGCTGGATTTTGCTTCATCACTAC				pcrsbe2con. seq
600	610	620	630	
ATTAAATACTTCTGAAGAGACAATTATTGATGA				10con. seq
ATTAAATACTTCTGAAGAGACAATTATTGATGA				11con. seq
ATTAAATACTTCTGAAGAGACAATTATTGATGA				19con. seq
ATTAAATACTTCTGAAGAGACAATTATTGATGA				86CON. SEQ
ATTAAATACTTCTGAAGAGACAATTATTGATGA				pcrsbe2con. seq
670	680	690	700	
GGACTTGGTCAGAAGATTTATGAAATAGACCCC				10con. seq
GGACTTGGTCAGAAGATTTATGAAATAGACCCC				11con. seq
GGACTTGGTCAGAAGATTTATGAAATAGACCCC				19con. seq
GGACTTGGTCAGAAGATTTATGAAATAGACCCC				86CON. SEQ
GGACTTGGTCAGAAGATTTATGAAATAGACCCC				pcrsbe2con. seq

Fig.10 SHEET 4

SUBSTITUTE SHEET (RULE 26)

50/75

	710	720	730
698	CTTTTGACAAACTATCGTCAACACCTTGATTACAGGT		
700	CTTTTGACAAACTATCGTCAACACCTTGATTACAGGT		
700	CTTTTGACAAACTATCGTCAACACCTTGATTACAGGT		
477	CTTTTGACAAACTATCGTCAACACCTTGATTACAGGT		
470	CTTTTGACAAACTATCGTCAACACCTTGATTACAGGT		
	780	790	800
768	ACAAGTATGAGGGTGGTTTGAAGCTTTTCTCGTGG		
770	ACAAGTATGAGGGTGGTTTGAAGC- TTTTCTCGTGG		
770	ACAAGTATGAGGGTGGTTTGAAGC TTTTCTCGTGG		
547	ACAAGTATGAGGGTGGTTTGAAGCTTTTCTCGTGG		
540	ACAAGTATGAGGGTGGTTTGAAGCTTTTCTCGTGG		
	850	860	870
838	AGGTATCACTTACCGTGAGTGGGCTCCTGGTGCCCAG		
839	AGGTATCACTTACCGTGAGTGGGCTCCTGGTGCCCAG		
840	AGGTATCACTTACCGTGAGTGGGCTC TGGTGCCCAG		
617	AGGTATCACTTACCGTGAGTGGGCTCCTGGTGCCCAG		
610	AGGTATCACTTACCGTGAGTGGGCTCCTGGTGCCCAG		
	920	930	940
908	GACGCAAATGCTGAC TTTATGACTCGGAATGAATTTG		
909	GACGCAAATGCTGACATTATGACTCGGAATGAATTTG		
910	GACGCAAATGCTGACATTATGACTCGGAATGAATTTG		
687	GACGCAAATGCTGACATTATGACTCGGAATGAATTTG		
680	GACGCAAATGCTGACATTATGACTCGGAATGAATTTG		
	990	1000	1010
978	ATGGTTCTCCTGCAATTCCTCATGGGTCCAGAGTGAA		
979	ATGGTTCTCCTGCAATTCCTCATGGGTCCAGAGTGAA		
980	ATGGTTCTCCTGCAATTCCTCATGGGTCCAGAGTGAA		
757	ATGGTTCTCCTGCAATTCCTCATGGGTCCAGAGTGAA		
750	ATGGTTCTCCTGCAATTCCTCATGGGTCCAGAGTGAA		

Fig.10  
Sheet 6

Fig.10 SHEET 5

SUBSTITUTE SHEET (RULE 26)

51/75

740	750	760	770	
ATTCACAGTACAAGAACTGAGGGAGGCAATTG				10con. seq
ATTCACAGTACAAGAACTGAGGGAGGCAATTG				11con. seq
ATTCACAGTACAAGAACTGAGGGAGGCAATTG				19con. seq
ATTCACAGTACAAGAACTGAGGGAGGCAATTG				86CON. SEQ
ATTCACAGTACAAGAACTGAGGGAGGCAATTG				pcrsbe2con. seq
810	820	830	840	
TTATGAAA <b>C</b> AATGGGTTTCACTCGTAGTGCTAC				10con. seq
TTATGAAAAAATGGGTTTCACTCGTAGTGCTAC				11con. seq
TTATGAAAAAATGGGTTTCACTCGTAGTGCTAC				19con. seq
TTATGAAAAAATGGGTTTCACTCGTAGTGCTAC				86CON. SEQ
TTATGAAAAAATGGGTTTCACTCGTAGTGCTAC				pcrsbe2con. seq
880	890	900	910	
TCAGCTGCCCTCATTGG <b>G</b> GATTTCACAATTGG				10con. seq
TCAGCTGCCCTCATTGGAGATTTCACAATTGG				11con. seq
TCAGCTGCCCTCATTGGAGATTTCACAATTGG				19con. seq
TCAGCTGCCCTCATTGGAGATTTCACAATTGG				86CON. SEQ
TCAGCTGCCCTCATTGGAGATTTCACAATTGG				pcrsbe2con. seq
950	960	970	980	
GTGTCTG <b>A</b> GAGATTTTTCTGCCAAATAATGTGG				10con. seq
GTGTCTGGGAGATTTTTCTGCCAAATAATGTGG				11con. seq
GTGTCTGGGAGATTTTTCTGCCAAATAATGTGG				19con. seq
GTGTCTGGGAGATTTTTCTGCCAAATAATGTGG				86CON. SEQ
GTGTCTGGGAGATTTTTCTGCCAAATAATGTGG				pcrsbe2con. seq
1020	1030	1040	1050	
GATACGTATGGACACTCCATCAGGTGTTAAGGA				10con. seq
GATACGTATGGACACTCCATCAGGTGTTAAGGA				11con. seq
GATACGTATGGACACTCCATCAGGTGTTAAGGA				19con. seq
GATACGTATGGACACTCCATCAGGTGTTAAGGA				86CON. SEQ
GATACG <b>T</b> ATGGACACTCCATCAGGTGTTAAGGA				pcrsbe2con. seq

Fig. 10 SHEET 6

52/75

	1060	1070	1080
1048	TTCCATTCCCTGCTTGGATCAACTACTCTTTACAGCTT		
1049	TTCCATTCCCTGCTTGGATCAACTACTCTTTACAGCTT		
1050	TTCCATTCCCTGCTTGGATCAACTACTCTTTACAGCTT		
827	TTCCATTCCCTGCTTGGATCAACTACTC--TACAGCTT		
820	TTCCATTCCCTGCTTGGATCAACTACTCTTTACAGCTT		
	1130	1140	1150
1118	GATCCACCCGAAGAGGAGAGGTATATCTTCCAACACC		
1119	GATCCACCCGAAGAGGAGAGGTATATCTTCCAACACC		
1120	GATCCACCCGAAGAGGAGAGGTATATCTTCCAACACC		
895	GATCCACCCGAAGAGGAGAGGTATATCTTCCAACACC		
890	GATCCACCCGAAGAGGAGAGGTATCTTCTTCCAACACC		
	1200	1210	1220
1188	ATGAATCTCATATTGGAATGAGTAGTCCGGAGCCTAA		
1189	ATGAATCTCATATTGGAATGAGTAGTCCGGAGCCTAA		
1190	ATGAATCTCATATTGGAATGAGTAGTCCGGAGCCTAA		
965	ATGAATCTCATATTGGAATGAGTAGTCCGGAGCCTAA		
960	ATGAATCTCATATTGGAATGAGTAGTCCGGAGCCTAA		
	1270	1280	1290
1258	TCTTCCTCGCATAAAAAAAGCTTGGGTACAATGCGCT		*
1259	TCTTCCTCGCATAAAAAA-GCTTGGGTACAATGCGCT		
1260	TCTTCCTCGCATAAAAAA-GCTTGGGTACAATGCGCT		
1035	TCTTCCTCGCATAAAAAA-GCTTGGGTACAATGCGCT		
1030	TCTTCCTCGCATAAAAAA-SCTTGGGTACAATGCGCT		
	1340	1350	1360
1328	TGCTAGTTTTGGTTATCATGTCACAAATTTTTTTTGCA		
1328	TGCTAGTTTTGGTTATCATGTCACAAATTTTTTTTGCA		
1329	CGCTAGTTTTGGTTATCATGTCACAAATTTTTTTTGCA		
1104	TGCTAGTTTTGGTTATCATGTCACAAATTTTTTTTGCA		
1099	TGCTAGTTTTGGTTATCATGTCACAAATTTTTTTTGCA		

Fig.10  
Sheet 8

Fig.10 SHEET 7

SUBSTITUTE SHEET (RULE 26)

53/75

1090	1100	1110	1120	
CCTGATGAAATTCCATATAATGGAATATATTAT	10con. seq			
CCTGATGAAATTCCATATAATGGAATATATTAT	11con. seq			
CCTGATGAAATTCCATATAATGGAATACATTAT	19con. seq			
CCTGATGAAATTCCATATAATGGAATATATTAT	86CON. SEQ			
CCTGATGAAATTCCATATAATGGAATATATTAT	pcrsbe2con. seq			
1160	1170	1180	1190	
CACGGCCAAAGAAACCAAAGTCGCTGAGAATAT	10con. seq			
CACGGCCAAAGAAACCAAAGTCGCTGAGAATAT	11con. seq			
CACGGCCAAAGAAACCAAAGTCGCTGAGAATAT	19con. seq			
CACGGCCAAAGAAACCAAAGTCGCTGAGAATAT	86CON. SEQ			
CACGGCCAAAGAAACCAAAGTCGCTGAGAATAT	pcrsbe2con. seq			
1230	1240	1250	1260	
AATTAACTCATACGTGAATTTTAGAGATGAAGT	10con. seq			
AATTAACTCATACGTGAATTTTAGAGATGAAGT	11con. seq			
AATTAACTCATACGTGAATTTTAGAGATGAAGT	19con. seq			
AATTAACTCATACGTGAATTTTAGAGATGAAGT	86CON. SEQ			
AATTAACTCATACGTGAATTTTAGAGATGAAGT	pcrsbe2con. seq			
1300	1310	1320	1330	
GCAAATTATGGCTATTCAAGAGCATTCTTATTA	10con. seq			
GC <del>G</del> AATTATGGCTATTCAAGAGCATTCTTATTA	11con. seq			
GCAAATTATGGCTATTCAAGAGCATTCTTATTA	19con. seq			
GCAAATTATGGCTATTCAAGAGCATTCTTATTA	86CON. SEQ			
GCAAATTATGGCTATTCAAGAGCATTCTTATTA	pcrsbe2con. seq			
1370	1380	1390	1400	
CCAAGCAGCCGTTTTTGGAAACGCCCGACGACCTT	10con. seq			
CCAAGCAGCCGTTTTTGGAAACGCCCGACGACCTT	11con. seq			
CCAAGCAGCCGTTTTTGGAAACGCCCGACGACCTT	19con. seq			
CCAAGCAGCCGTTTTTGGAAACGCCCGACGACCTT	86CON. SEQ			
CCAAGCAGCCGTTTTTGGAAACGCCCGACGACCTT	pcrsbe2con. seq			

Fig. 10 SHEET 8

SUBSTITUTE SHEET (RULE 26)

54/75

	1410	1420	1430
1398	AAGTCTTTGATTGATAAAGCTCATGAGCTAGGAATTG		
1398	AAGTCTTTGATTGATAAAGCTCATGAGCTAGGAATTG		
1399	AAGTCTTTGATTGATAAAGCTCATGAGCTAGGAATTG		
1174	AAGTCTTTGATTGATAAAGCTCATGAGCTAGGAATTG		
1169	AAGTCTTTGATTGATAAAGCTCATGAGCTAGGAATTG		
	1480	1490	1500
1468	CAAATAATACTTTAGATGGACTGAACATGTTTGACGG		
1468	CAAATAATACTTTAGATGGACTGAACATGTTTGACGG		
1469	CAAATAATACTTTAGATGGACTGAACATGTTTGACGG		
1244	CAAATAATACTTTAGATGGACTGAACATGTTTGACGG		
1239	CAAATAATACTTTAGATGGACTGAACATGTTTGACGG		
	1550	1560	1570
1538	TGGTTATCATTGGATGTGGGATTCCGCCTCTTTAAC		
1538	TGGTTATCATTGGATGTGGGATTCCGCCTCTTTAAC		
1539	TGGTTATCATTGGATGTGGGATTCCGCCTCTTTAAC		
1314	TGGTTATCATTGGATGTGGGATTCCGCCTCTTTAAC		
1309	TGGTTATCATTGGATGTGGGATTCCGCCTCTTTAAC		
	1620	1630	1640
1608	TCAAATGCGAGATGGTGGTTGGATGAGTTCAAATTTG		
1607	TCAAATGCGAGATGGTGGTTGGATGAGTTCAAATTTG		
1609	TCAAATGCGAGATGGTGGTTGGATGAGTTCAAATTTG		
1384	TCAAATGCGAGATGGTGGTTGGATGAGTTCAAATTTG		
1379	TCAAATGCGAGATGGTGGTTGGATGAGTTCAAATTTG		
	1690	1700	1710
1678	TGTACTCACCACGGATTATCGGTGGGATTCACTGG		
1677	TGTATACTCACCACGGATTATCGGTGGGATTCACTGG		
1679	TGTATACTCACCACGGATTATCGGTGGGATTCACTGG		
1454	TGTATACTCACCACGGATTATCGGTGGGATTCACTGG		
1449	TGTATACTCACCACGGATTATCGGTGGGATTCACTGG		

Fig. 10  
Sheet 10

Fig. 10 SHEET 9

SUBSTITUTE SHEET (RULE 26)



55/75

1440	1450	1460	1470	
TTGTTCTCATGGACATTGTTT	CACAGCCATGCAT	10con. seq		
TTGTTCTCATGGACATTGTTT	CACAGCCATGCAT	11con. seq		
TTGTTCTCATGGACATTGTTT	CACAGCCATGCAT	19con. seq		
TTGTTCTCATGGACATTGTTT	CACAGCCATGCAT	86CON. SEQ		
TTGTTCTCATGGACATTGTTT	CACAGCCATGCAT	pcrsbe2con. seq		
1510	1520	1530	1540	
CACAGATAGTTGTTACTTTT	CACTCTGGAGCTCG	10con. seq		
CACCGATAGTTGTTACTTTT	CACTCTGGAGCTCG	11con. seq		
CACCGATAGTTGTTACTTTT	CACTCTGGAGCTCG	19con. seq		
CACCGATAGTTGTTACTTTT	CACTCTGGAGCTCG	86CON. SEQ		
CACAGATAGTTGTTACTTTT	CACTCTGGAGCTCG	pcrsbe2con. seq		
1580	1590	1600	1610	
TATGGAAACTGGGAGGTACTT	AGGTATCTTCTC	10con. seq		
TATGGAAACTGGGAGGTACTT	AGGTATCTTCTC	11con. seq		
TATGGAAACTGGGAGGTACTT	AGGTATCTTCTC	19con. seq		
TATGGAAACTGGGAGGTACTT	AGGTATCTTCTC	86CON. SEQ		
TATGGAAACTGGGAGGTACTT	AGGTATCTTCTC	pcrsbe2con. seq		
1650	1660	1670	1680	
ATGGATTTAGATTTGATGGT	GTGACATCAATGA	10con. seq		
ATGGATTTAGATTTGATGGT	GTGACATCAATGA	11con. seq		
ATGGATTTAGATTTGATGGT	GTGACATCAATGA	19con. seq		
ATGGATTTAGATTTGATGGT	GTGACATCAATGA	86CON. SEQ		
ATGGATTTAGATTTGATGGT	GTGACATCAATGA	pcrsbe2con. seq		
1720	1730	1740	1750	
GAACTACGAGGAATACTTTT	GGACTCGCAACTGA	10con. seq		
GAACTACGAGGAATACTTTT	GGACTCGCAACTGA	11con. seq		
GAACTACGAGGAATACTTTT	GGACTCGCAACTGA	19con. seq		
GAACTACGAGGAATACTTTT	GGACTCGCAACTGA	86CON. SEQ		
GAACTACGAGGAATACTTTT	GGACTCGCAACTGA	pcrsbe2con. seq		

Fig. 10 SHEET 10

SUBSTITUTE SHEET (RULE 26)

56/75

	1760	1770	1780
1748	TGTGGATGCTGTTGTGTATCTGATGCTGGTCAACGAT		
1747	TGTGGATGCTGTTGTGTATCTGATGCTGGTCAACGAT		
1749	TGTGGATGCTGTTGTGTATCTGATGCTGGTCAACGAT		
1524	TGTGGATGCTGTTGTGTATCTGATGCTGGTCAACGAT		
1519	TGTGGATGCTGTTGTGTATCTGATGCTGGTCAACGAT		
	1830	1840	1850
1818	ATTGGTGAAGATGTTAGCGGAATGCCGACATTTTGTG		
1817	ATTGGTGAAGATGTTAGCGGAATGCCGACATTTTGTG		
1819	ATTGGTGAAGATGTTAGCGGAATGCCGACATTTTGTG		
1594	ATTGGTGAAGATGTTAGCGGAATGCCGACATTTTGTG		
1589	ATTGGTGAAGATGTTAGCGGAATGCCGACATTTTGTG		
	1900	1910	1920
1888	ATCGGCTGCATATGGCAATTGCTGATAAATGGATTGA		
1887	ATCGGCTGCATATGGCAATTGCTGATAAATGGATTGA		
1889	ATCGGCTGCATATGGCAATTGCTGATAAATGGATTGA		
1664	ATCGGCTGCATATGGCAATTGCTGATAAATGGATTGA		
1659	ATCGGCTGCATATGGCAATTGCTGATAAATGGATTGA		
	1970	1980	1990
1958	GGGTGATATTGTTTCATACACTGACAAATAGAAGATGG		
1957	GGGTGATATTGTTTCATACACTGACAAATAGAAGATGG		
1959	GGGTGATATTGTTTCATACACTGACAAATAGAAGATGG		
1734	GGGTGATATTGTTTCATACACTGACAAATAGAAGATGG		
1729	GGGTGATATTGTTTCATACACTGACAAATAGAAGATGG		
	2040	2050	2060
2028	GATCAAGCTCTAGTCGGTGATAAACTATAGCATTCT		
2027	GATCAAGCTCTAGTCGGTGATAAACTATAGCATTCT		
2029	GATCAAGCTCTAGTCGGTGATAAACTATAGCATTCT		
1804	GATCAAGCTCTAGTCGGTGATAAACTATAGCATTCT		
1799	GATCAAGCTCTAGTCGGTGATAAACTATAGCATTCT		

Fig. 10  
Sheet 12

Fig. 10 SHEET 11

SUBSTITUTE SHEET (RULE 26)

57/75

1790	1800	1810	1820	
CTTATTTCATGGGCTTTTCCCAGATGCAATTACC	10con. seq			
CTTATTTCATAGGCTTTTCCCAGATGCAATTACC	11con. seq			
CTTATTTCATGGGCTTTTCCCAGATGCAATTACC	19con. seq			
CTTATTTCATGGGCTTTTCCCAGATGCAATTACC	86CON. SEQ			
CTTATTCAAGGCTTTTCCCAGATGCAATTACC	pcrsbe2con. seq			
1860	1870	1880	1890	
TTCCCGTTCAAGATGGGGGTGTTGGCTTTGACT	10con. seq			
TTCCCGTTCAAGATGGGGGTGTTGGCTTTGACT	11con. seq			
TTCCCGTCAAGAAGGGGGGTGTTGGCTTTGACT	19con. seq			
TTCCCGTTCAAGATGGGGGTGTTGGCTTTGACT	86CON. SEQ			
TTCCCGTTCAAGATGGGGGTGTTGGCTTTGACT	pcrsbe2con. seq			
1930	1940	1950	1960	
GTTGCTCAAGAAACGGGATGAGGATTGGAGAGT	10con. seq			
GTTGCTCAAGAAACGGGATGAGGATTGGAGAGT	11con. seq			
GTTGCTCAAGAAACGGGATGAGGATTGGAGAGT	19con. seq			
GTTGCTCAAGAAACGGGATGAGGATTGGAGAGT	86CON. SEQ			
GTTGCTCAAGAAACGGGATGAGGATTGGAGAGT	pcrsbe2con. seq			
2000	2010	2020	2030	
TCGGAAAAGTGTGTTTCATACGCTGAAAGTCAT	10con. seq			
TCGGAAAAGTGTGTTTCATACGCTGAAAGTCAT	11con. seq			
TCGGAAAAGTGTGTTTCATACGCTGAAAGTCAT	19con. seq			
TCGGAAAAGTGTGTTTCATACGCTGAAAGTCAT	86CON. SEQ			
TCGGAAAAGTGTGTTTCATACGCTGAAAGTCAT	pcrsbe2con. seq			
2070	2080	2090	2100	
GGCTGATGGACAAGGATATGTATGATTTTATGG	10con. seq			
GGCTGATGGACAAGGATATGTATGATTTTATGG	11con. seq			
GGCTGATGGACAAGGATATGTATGATTTTATGG	19con. seq			
GGCTGATGGACAAGGATATGTATGATTTTATGG	86CON. SEQ			
GGCTGATGGACAAGGATATGTATGATTTTATGG	pcrsbe2con. seq			

Fig. 10 SHEET 12

SUBSTITUTE SHEET (RULE 26)

58/75

	2110	2120	2130
2098	CTCTGGATAGACCGT	CAACATCATT	AATAGATCGTGG
2097	CTCTGGATAGACCG	CAACATCATT	AATAGATCGTGG
2099	CTCTGGATAGACCGT	CAACATCATT	AATAGATCGTGG
1874	CTCTGGATAGACCG	CAACATCATT	AATAGATCGTGG
1869	CTCTGGATAGACCG	CAACATCATT	AATAGATCGTGG
	2180	2190	2200
2168	TATGGGATTAGGAGG	AAGGGTACCT	AAATTTTCATG
2167	TATGGGATTAGGAGG	AAGGGTACCT	AAATTTTCATG
2169	TATGGGATTAGGAGG	AAGGGTACCT	AAATTTTCATG
1944	TATGGGATTAGGAGG	AAGGGTACCT	AAATTTTCATG
1939	TATGGGATTAGGAGG	AAGGGTACCT	AAATTTTCATG
	2250	2260	2270
2238	TTCCCTAGGGCTGA	ACAACACCT	TCTCTGATGGCTCAG
2237	TTCCCTAGGGCTGA	ACAACACCT	TCTCTGATGGCTCAG
2239	TTCCCTAGGGCTGA	ACAACACCT	TCTCTGATGGCTCAG
2014	TTCCCTAGGGCTGA	ACAACACCT	TCTCTGATGGCTCAG
2009	TTCCCTAGGGCTGA	ACAACACCT	TCTCTGATGGCTCAG
	2320	2330	2340
2308	GCAGACGGAGATTT	GACCTGGGAGAT	GCAGAATATTT
2307	GCAGACGGAGATTT	GACCTGGGAGAT	GCAGAATATTT
2309	GCAGACGGAGATTT	GACCTGGGAGAT	GCAGAATATTT
2084	GCAGACGGAGATTT	GACCTGGGAGAT	GCAGAATATTT
2079	GCAGACGGAGATTT	GACCTGGGAGAT	GCAGAATATTT
	2390	2400	2410
2378	TATGCAGTATCTT	GAAGATAAAT	TATGAGTTTATGACT
2377	TATGCAGTATCTT	GAAGATAAAT	TATGAGTTTATGACT
2379	TATGCAGTATCTT	GAAGATAAAT	TATGAGTTTATGACT
2154	TATGCAGTATCTT	GAAGATAAAT	TATGAGTTTATGACT
2149	TATGCAGTATCTT	GAAGATAAAT	TATGAGTTTATGACT

Fig.10  
Sheet 14

Fig. 10 SHEET 13

SUBSTITUTE SHEET (RULE 26)

59/75

2140	2150	2160	2170	
GATAGCATT	ACACAAGATGATTAGGCTTGTAAC			10con. seq
GATAGCATT	GCACAAGATGATTAGGCTTGTAAC			11con. seq
GATAGCATT	GCACAAGATGATTAGGCTTGTAAC			19con. seq
GATAGCATT	GCACAAGATGATTAGGCTTGTAAC			86CON. SEQ
GATAGCATT	GCACAAGATGATTAGGCTTGTAAC			pcrsbe2con. seq

2210	2220	2230	2240	
GGAAATGAATT	CGGCCACCCTGAGTGGATTGAT			10con. seq
GGAAATGAATT	CGGCCACCCTGAGTGGATTGAT			11con. seq
GGAAATGAATT	CGGCCACCCTGAGTGGATTGAT			19con. seq
GGAAATGAATT	CGGCCACCCTGAGTGGATTGAT			86CON. SEQ
GGAAATGAATT	CGGCCACCCTGAGTGGATTGAT			pcrsbe2con. seq

2280	2290	2300	2310	
TAATTCCC	AGAAACCAATTCAGTTATGATAAAT			10con. seq
TAATTCCC	GGAAACCAATTCAGTTATGATAAAT			11con. seq
TAAT	CGCGAAACCAATTCAGTTATGATAAAT			19con. seq
TAATTCCC	GGAAACCAATTCAGTTATGATAAAT			86CON. SEQ
TAATTCCC	GGAAACCAATTCAGTTATGATAAAT			pcrsbe2con. seq

2350	2360	2370	2380	
AAGATACCGTGGGTTGCAAGAATTTGACCGGGC				10con. seq
AAGATACC	ATGGGTTACAAGAATTTGAC	TGGGC		11con. seq
AAGATACCGTGGGTTGCAAGAATTTGACCGGGC				19con. seq
AAGATACCGTGGGTTGCAAGAATTTGACCGGGC				86CON. SEQ
AAGATACC	ATGGGTTGCAAGAATTTGACCGGGC			pcrsbe2con. seq

2420	2430	2440	2450	
TCAGAACACCAAGTTTCATATCACGAAAGGATGAA				10con. seq
TCAGAACACCAAGTTTCATATCACGAAAGGATGAA				11con. seq
TCAGAACACCAAGTTTCATATCACGAAAGGATGAA				19con. seq
TCAGAACACCAAGTTTCATATCACGAAAGGATGAA				86CON. SEQ
TCAGAACACCAAGTTTCATATCACGAAAGGATGAA				pcrsbe2con. seq

Fig. 10 SHEET 14

SUBSTITUTE SHEET (RULE 26)

60/75

	2460	2470	*	2480
2448	GGAGATAGGATGATTGTATTTGAAAAAGGAAACCTAG			
2447	GGAGATAGGATGATTGTATTTGAAAAGAGGAAACCTAG			
2449	GGAGATAGGATGATTGTATTTGAAAAAGGAAACCTAG			
2224	GGAGATAGGATGATTGTATTTGAAAAAGGAAACCTAG			
2219	GGAGATAGGATGATTGTATTTGAAAAGAGGAAACCTAG			
			*	
	2530	2540		2550
2518	ATTCAGACTATCGCATAGGCTGCCTGAAGCCTGGAAA			
2517	ATTCAGACTATCGCATAGGCTGCCTGAAGCCTGGAAA			
2519	ATTCAGACTATCGCATAGGCTGCCTGAAGCCTGGAAA			
2294	ATTCAGACTATCGCATAGGCTGCCTGAAGCCTGGAAA			
2289	ATTCAGACTATCGCATAGGCTGCCTGAAGCCTGGAAA			
	2600	2610		2620
2588	TTTTGGTGGCTTCGGGAGAATTGATCATAATGCCGAA			
2587	TTTTGGTGGCTTCGGGAGAATTGATCATAATGCCGAA			
2589	TTTTGGTGGCTTCGGGAGAATTGATCATAATGCCGAA			
2364	TTTTGGTGGCTTCGGGAGAATTGATCATAATGCCGAA			
2359	TTTTGGTGGCTTCGGGAGAATTGATCATAATGCCGAA			
	2670	2680	*	2690
2658	CCTCGTTCAATTATGGTGTATGCACCTAGTAGAACAG			
2657	CCTCGTTCAATTATGGTGTATGCACCTAGTAGAACAG			
2659	CCTCGTTCAATTATGGTGTATGCACCTAGTAGAACAG			
2434	CCTCGTTCAATTATGGTGTATGCACCTAGTAGAACAG			
2429	CCTCGTTCAATTATGGTGTATGCACCTAGTAGAACAG			
			*	
	2740	2750		2760
2722	-----AAGAAGAAGAAGAAGAAGAAGTAGCAGTAGT			
2722	-----AGAAGTAGCAGTAGT			
2729	AAGAAGAAGAAGAAGAAGAAGAAGAAGTAGCAGTAGT			
2501	AAGAAGAAGAAGAAGAAGAAGAAGAAGTAGCAGTAGT			
2499	NAGAAGAAGAAGAAGAN-----			

Fig. 10  
Sheet 16

Fig. 10 SHEET 15

SUBSTITUTE SHEET (RULE 26)

61/75

2490	2500	2510	*	2520	
TTTTTGTCTTTAATTTTCACTGGACAAAAGGCT					10con. seq
TTTTTGTCTTTAATTTTCACTGGACAAAAGCT					11con. seq
TTTTTGTCTTTAATTTTCACTGGACAAAAGCT					19con. seq
TTTTTGTCTTTAATTTTCACTGGACAAAAGCT					86CON. SEQ
TTTTTGTCTTTAATTTTCACTGGACAAAAGCT					pcrsbe2con. seq
					*
2560	2570	2580		2590	
ATACAAGGTTGCCTTGGACTCAGATGATCCACT					10con. seq
ATACAAGGTTGCTTGGACTCAGATGATCCACT					11con. seq
ATACAAGGTTGCCTTGGACTCAGATGATCCACT					19con. seq
ATACAAGGTTGCCTTGGACTCAGATGATCCACT					86CON. SEQ
ATACAAGGTTGCTTGGACTCAGATGATCCACT					pcrsbe2con. seq
					*
2630	*	2640	*	2650	2660
TATTTACCTTTGAAGGATGGTATGATGATCGT					10con. seq
TATTTACCTTTGAAGGATGGTATGATGATCGT					11con. seq
TATTTACCTTTGAAGGATGGTATGATGATCGT					19con. seq
TATTTACCTTTGAAGGATGGTATGATGATCGT					86CON. SEQ
TATTTACCTTTGAAGGATGGTATGATGATCGT					pcrsbe2con. seq
					*
2700	2710	2720		2730	
CAGTGGTCTATGCACTAGTAGACAAAG---					10con. seq
CAGTGGTCTATGCACTAGTAGACAAAGCT---					11con. seq
CAGTGGTCTATGCACTAGTAGACAAAGAAAGAAG					19con. seq
CAGTGGTCTATGCACTAGTAGACAAAG---AAG					86CON. SEQ
CAGTGGTCTATGCACTAGTAGACAAAGNTAGAAG					pcrsbe2con. seq
					*
2770	2780	2790		2800	
AGAAGAAGTAGTAGTAGAAGAAGAATGAACGAA					10con. seq
AGAAGAAAGCCCATTTG-----AAGAATGAACGAA					11con. seq
AGAAGAAGTAGTAGTAGAAGAAGAATGAACGAA					19con. seq
AGAAGAAGTAGTAGTAGAAGAAGAATGAACGAA					86CON. SEQ
-----CCGNNGAAGAAT-----					pcrsbe2con. seq

Fig. 10 SHEET 16

SUBSTITUTE SHEET (RULE 26)

62/75

	2810	2820	2830
2786	CTTGTGATCGCGTTGAAAGATTTGAACGCC	CACATAGA	
2764	CTTGTGATCGCGTTGAAAGATTTGAACG	TAC	TGG-
2799	CTTGTGATCGCGTTGAAAGATTTGAACG	CTACATAGA	
2571	CTTGTG		
2529			
	2880	2890	2900
2856	CTTGGCGGAATTT	CATGTGACAACA	-GGTTTGCAATT
2829	CTTGGCGGAATTT	CATGTGACAACA	AGGTTTGCA
2869	CTTGGCGGAATTT	CATGTGACA	CAA-GGTTTGCAATT
2576			
2529			
	2950	2960	2970
2925	GAGATGAAGTGCTGAACAAA	AA	CATATGTAAAATCGA
2899	GAGATGAAGTGCTGAACAAA	--	CATATGTAAAATCGA
2938	GAGATGAAGTGCTGAACAAA	--	CATATGTAAAATCGA
2576			
2529			
	3020	3030	
2995	CCTGCAG		CC
2967	CCTGCAG		CC
3006	CCTGCAG	GCCGGGGGACCCCTTAGTT	CT
2576			
2529			


Fig.10  
Sheet 18


Fig. 10 SHEET 17


SUBSTITUTE SHEET (RULE 26)



63/75

2840	2850	2860	2870	
GCTTCTTGACGTATCTGGCAATATTGCATTAGT				10con. seq
--TCATCCACATA--GAGCTTCTTGACATCAGT				11con. seq
GCTTCTTGACGTATCTGGCAATATTGCATCAGT				19con. seq
				86CON. SEQ
-----				pcrsbe2con. seq

2910	2920	2930	2940	
CTTTCCACTATTAGTAGTGCAACGATATACGCA				10con. seq
CTTTCCACTATTAGTAGTCCACCGATATACGCA				11con. seq
CTTTCCACTATTAGTAGTGCAACGATATACGCA				19con. seq
				86CON. SEQ
-----				pcrsbe2con. seq

2980	2990	3000	3010	
TGAATTTATGTCGAATGCTGGGACGATCGAATT				10con. seq
TGAATTTATGTCGAATGCTGGGACGATCGAATT				11con. seq
TGAATTTATGTCGAATGCTGGGACGATCGAATT				19con. seq
				86CON. SEQ
-----				pcrsbe2con. seq

				10con. seq
				11con. seq
				19con. seq
				86CON. SEQ
				pcrsbe2con. seq

Fig. 10 SHEET 18

SUBSTITUTE SHEET (RULE 26)

64/75




Fig. 11

65/75

Nco I  
 BstX I  
 TCATTAAGAGGAGAAATTAACTATGAGAGGATCTCACCATCACCATCACCATGGGATCT 60  
 AGTAATTTCTCCTCTTAATGATACTCTCCTAGAGTGGTAGTGGTAGCCCTAGA  
 M R G S H H H H H G I

EcoR I  
 TGGCTGAAAAGTCITTTACAATTCCGAATTCGACCTTCTACAGTTGCAGCATCGGGGA 120  
 ACCGACTTTTCAGAAGAATGTTAAGGCTTAAGGCTGGAAGATGTCAACGTCGTAGCCCT  
 L A E K S S Y N S E F R P S T V A A S G  
 AAGTCCTTGTGCCTGGAACCCAGAGTGATAGCTCCTCATCCTCAACAAACCAATTTGAGT 180  
 TTCAGGAACACGGACCTTGGGTCTCACTATCGAGGAGTAGGAGTTGTTTGGTTAAACTCA  
 K V L V P G T Q S D S S S S S T N Q F E

TCACTGAGACATCTCCAGAAAATTCGCCAGCATCAACTGATGTAGATAGTTCACCAATGG 240  
 AGTGACTCTGTAGAGGTCITTTAAGGGTCGTAGTTGACTACATCTATCAAGTTGTTACC  
 F T E T S P E N S P A S T D V D S S T M

Fig.12  
SHEET 1

66/75

AACACGCTAGCCAGATTAAACTGAGAACGATGACGTTGAGCCGTCAAGTGATCTTACAG  
TTGTGCGATCGGTCATAATTIGACTCTTGCTACTGCAACTCGGCAGTTCAC TAGAATGTC  
E H A S Q I K T E N D D V E P S S D L T

GAAGTGTTGAAGAGCTGGATTTTGCTTCATCACTACAAGAGGTGGTAAACTGG  
CTTCACAACTTCGACCTAAACGAAGTAGTGATGTTGATGTTCTTCCACCATTTGACC  
G S V E E L D F A S S L Q L Q E G G K L

AGGAGTCTAAACATTAAATACTTCTGAAGAGACAATTATGATGAATCTGATAGGATCA  
TCCTCAGATTTTGTAATTTATGAAGACTTCTCTGTTAATACTACTTAGACTATCCTAGT  
E E S K T L N T S E E T I I D E S D R I

GAGAGAGGGGCATCCCTCCACCTGGACTTGGTCAGAAGATTATGAATAGACCCCTTT  
CTCTCTCCCGTAGGGAGGTGGACCTGAACCACTTCTTAATACTTTATCTGGGGGAAA  
R E R G I P P P G L G Q K I Y E I D P L

Hinc II

TGACAAACTATCGTCAACACCTTGATTACAGGTATTCACAGTACAAGAACTGAGGGAGG  
ACTGTTTGATAGCAGTTGTGGAACATAATGTCCATAAGTGTCATGTTCTTTGACTCCCTCC  
L T N Y R Q H L D Y R Y S Q Y K K L R E

Fig.12  
SHEET 2

SUBSTITUTE SHEET (RULE 26)

67/75

Hind III  
CAATIGACAAGTATGAGGGTGGTTTGGAGCTTTTCTCGTGGTTATGAAAAATGGGT  
GTTAACTGTTTACTCTCCACCAACCTTCGAAAAAGAGCACCATACTTTTACCCAA  
A I D K Y E G G L E A F S R G Y E K M G 600

Pvu II  
TCACTCGTAGTGCTACAGGTATCATTACCGTGAGTGGCTCCTGGTGCCAGTCAGCTG  
AGTGAGCATCAGGATGTCCTAGTGAATGGCAGTCCCGAGGACCACGGGTCAGTCGAC  
F T R S A T G I T Y R E W A P G A Q S A 660

CCCICATTGGAGATTTCACAAATTGGGACGCAATGCTGACATTATGACTCGGAATGAAT  
GGGAGTAACCTCTAAAGTTGTTAACCCTGGTTACGACTGTAATACTGAGCCTTACTTA  
A L I G D F N N W D A N A D I M T R N E 720

TTGGTGTCTGGGAGATTTTCTGCCAAATAATGTGGATGGTTCTCCTGCAATTCCTCATG  
AACCACAGACCCCTCTAAAAGACGGTTTATTACACCTACCAAGAGGACGTTAAGGAGTAC  
F G V W E I F L P N N V D G S P A I P H 780

Fig 12  
SHEET 3

68/75

SnaBI

GGTCCAGAGTGAAGATACGTATGGACACTCCATCAGGTGTTAAGGATTCATTCCITGCTT 840

CCAGGTCACATTCATGCATACCTGTGAGGTAGTCCACAATTCCTAAGGTAAGGACGAA

G S R V K I R M D T P S G V K D S I P A

GGATCAACTACTCTTCACAGCTTCCCTGATGAAATCCATATAATGGAATATATATGATC 900

CCTAGTTGATGAGAAGTGTGGAAGGACTACTTTAAGGTATATTACCTTATATAATACTAG

W I N Y S S Q L P D E I P Y N G I Y Y D

CACCCGAAGAGGAGGTATATCTTCCAACACCCAGGCCAAAGAAACCAAGTCGCTGA 960

GTGGGCTTCTCCTCCATATAGAAGTTGTGGTGCCGGTTCTTTGGTTTCAGCGACT

P P E E R Y I F Q H P R P K K P K S L

GAATATGAATCTCATATTGGAATGAGTAGTCCGGAGCCTAAAATTAACATCATACGIGA 1020


CTTATATACTTAGAGTATAACCTTACTCATCATCAGGCCTCGGATTTTAAATTGAGTATGCACT

R I Y E S H I G M S S P E P K I N S Y V

SUBSTITUTE SHEET (RULE 26)

Fig. 12  
SHEET 4

69/75

Fig. 12  
SHEET 5

70/75

SacI

ATGGACTGAACATGTTTGACGGCACCAGTAGTTGTTACTTTCACTCIGGAGCTCGTGGTT  
TACCTGACTTGTACAAACTGCCGTGGCTATCAACAATGAAAGTGAGACCTCGAGCACCAA  
D G L N M F D D G T D S C Y F H S G A R G 1320

ATCATTGGATGTGGGATTCGCCCTTTTAACTATGGAACCTGGGAGGTACTTAGGTATC  
TAGTAACCTACACCCCTAAGGGCGGAAATTTGATACCTTTGACCCCTCCATGAATCCATAG  
Y H W M W D S R L F N Y G N W E V L R Y 1380

TTCTCTCAAATGCGAGATGGTGGTTGGATGAGTTCAAATTTGATGGATTTAGATTGATG  
AAGAGAGTTTACGCTCTACCCACCACTACTCAAGTTTAACTACCTAAATCTAAACTAC  
L L S N A R W W L D E F K F D G F R F D 1440

GTGTGACATCAATGATGTATACACACCAGGATTATCGGTGGGATTCACCTGGGAACACG  
CACACTGTAGTTACTACATATGAGTGGTGCCTAATAGCCACCCCTAAGTGACCCCTTGATGC  
G V T S M M Y T H H G L S V G F T G N Y 1500

SUBSTITUTE SHEET (RULE 26)

Fig. 12  
SHEET 6



71/75

Hinc II

AGGAATACITTTGGACITCGCAACTGATGIGGATGCTGTTGIGTATCTGATGCTGGTCAACG 1560

TCCTTATGAAACCTGAGCGTTGACTACACCTACGACAACACATAGACTACGACCAGTTGC  
E E Y F G L A T D V D A V V Y L M L V N

ATCTTATCATGGGCTTTTCCAGATGCAATTACCATTTGGTGAAGATGTTAGCGGAATGC 1620

TAGAATAAGTACCCGAAAGGCTCTACGTTAATGGTAACCACTTCTACAATCGCCTTACG  
D L I H G L F P D A I T I G E D V S G M

CGACATTTTGATTCGGTTCAAGATGGGGTGTGGCTTTGACTATCGGCTGCATATGG 1680

GCTGTAAACATAAGGGCAAGTTCTACCCCAACCGAACTGATAGCCGACGTATACC  
P T F C I P V Q D G G V G F D Y R L H M

CAATTGCTGATAAATGGATTGAGTTGCTCAAGAAACGGGATGAGGATTGGAGAGTGGGTG 1740

GTTAACGACTATTTACCTAACTCAACGAGTTCTTTGCCCTACTCCTAACCTCTCACCCAC  
A I A D K W I E L L K K R D E D W R V G

ATATIGTTTCATACACTGACAAATAGAAGATGGTCGGAAGAGTGIGTTTCATACGCTGAAA 1800

TATAACAAGTATGTGACTGTTTATCTTCTACCGCCTTTTCACACAAGTATGCGACTTT  
D I V H T L T N R R W S E K C V S Y A E

SUBSTITUTE SHEET (RULE 26)

Fig 12  
SHEET 7

72/75

GTCATGATCAAGCTCTAGTCGGTGATAAACTATAGCATTCTGGCTGATGGACAAGGATA 1860  
 CAGTACTAGTTCGAGATCAGCCACTATTTTGATATCGTAAGACCGACTACCTGTTCCAT  
 S H D Q A L V G D K T I A F W L M D K D

TGATGATTTTATGGCTCIGGATAGACCGCCAACATCATTAATAGATCGTGGGATAGCAT 1920  
 ACATACTAAATACCGAGACCTATCTGGCGTTGTAGTAATTAATCTAGCACCCCTATCGTA  
 M Y D F M A L D R P P T S L I D R G I A

Asp 718

Kpn I

TGCACAAGATGATTAGGCTTGTAACATATGGGATTAGGAGGAGGAGGTACCTAAATTICA 1980  
 ACGTGTCTACTAATCCGAACATTGATACCCCTAATCCCTCCTCTCCCATGGATTAAAGT  
 L H K M I R L V T M G L G G E G Y L N F

EcoRI

TGGGAAATGAATTCGGCCACCCIGAGTGGATTGATTCCCTAGGGCTGAACAACACCTCT 2040  
 ACCCTTTACTTAAGCCGGTGGGACTCACCTAACTAAGGGATCCCGACTTGTGTGGAGA  
 M G N E F G H P E W I D F P R A E Q H L

Fig 12  
SHEET 8

SUBSTITUTE SHEET (RULE 26)

73/75

CTGATGACTCAGTAATCCCGGAAACCAATTCAGTTATGATAAATGCAGACGGAGATTG 2100  
GACTACTGAGTCATTAAGGCCCTTTGGTTAAGTCAATACTATTTACGTCTGCCCTCTAAAC  
S D D S V I P G N Q F S Y D K C R R R F

Ssp I

ACCTGGGAGATGCAGAATAATTAAGATACCGTGGTTGCAAGAAATTTGACCGGGCTATGC 2160  
TGGACCCCTCAGTCTTATAAATTCATGGCACCCCAACGTTCTTAACTGGCCCGGATACG  
D L G D A E Y L R Y R G L Q E F D R A M

AGTATCTTGAAGATAAATAGTATTATGACTTCAGAACACCAGTTCATATCAGAAAGG 2220  
TCATAGAACTTCTATTATCTCAATACTGAAGTCTTGTGGTCAAGTATAGTCTTTCC  
Q Y L E D K Y E F M T S E H Q F I S R K

ATGAAGGAGATAGGATGATTGTATTGAAAAAGGAAACCTAGTTTTTGTCITTAATTTT 2280  
TACTTCCTCTATCCTACTAACATAAACTTTTTCCTTTGGATCAAAAACAGAAATTTAAAG  
D E G D R M I V F E K G N L V F V F N F

ACTGGACAAAAGCTATTTCAGACTATCGCATAGGCTGCCTGAAGCCTGGAAAATACAAG 2340  
TGACCTGTTTTTCGATAAGTCTGATAGCGTATCCGACGGACTTCGGACCTTTTATGTTCC  
H W T K S Y S D Y R I G C L K P G K Y K

SUBSTITUTE SHEET (RULE 26)

Fig. 12  
SHEET 9



75/75




Fig. 13

SUBSTITUTE SHEET (RULE 26)

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**